

ON THE GEOMETRY OF A CERTAIN SLICE OF THE CHARACTER VARIETY OF A KNOT GROUP

FUMIKAZU NAGASATO AND YOSHIKAZU YAMAGUCHI

ABSTRACT. We construct explicitly a map from a certain slice (an algebraic subset) of the $\mathrm{SL}_2(\mathbb{C})$ -character variety of a knot exterior to the $\mathrm{SL}_2(\mathbb{C})$ -character variety of a two-fold branched cover along the knot. This map has a property such as a covering map for a two-fold branched cover. Namely, if the map is surjective, the slice has a two-fold branched cover-like structure whose base space can be thought of as the $\mathrm{SL}_2(\mathbb{C})$ -character variety of the two-fold branched cover and whose branched point set consists entirely of the characters of metabelian representations of the knot group. By using this map, we show that all abelian characters for a two-fold branched cover along a knot are given by the characters of metabelian representations of the knot group and its representatives can be taken by binary dihedral representations of the knot group. Moreover, we observe that the above map is surjective for two-bridge knots and pretzel knots of type (p, q, r) . The above framework can be naturally extended to more general setting (i.e., n -fold cyclic branched covers).

1. INTRODUCTION

In low-dimensional topology and geometry, the character variety of the fundamental group of a manifold and its associated topological invariants have been playing important roles. The G -character variety of a manifold M can be considered as the set of characters of G -representations of the fundamental group $\pi_1(M)$ in a sense (for a precise definition for $\mathrm{SL}_2(\mathbb{C})$, refer to Subsection 2.2 or the original article [CS83]). To talk about the history on invariants associated to the character varieties, we first recall the case of $\mathrm{SU}(2)$. For an integral homology three-sphere Σ , the Casson invariant is defined originally by using the intersection between $\mathrm{SU}(2)$ -character varieties associated to a Heegaard splitting of Σ (see [AM90, Sav99] for expositions). For a knot in three-sphere S^3 , X-S. Lin [Lin92] has defined so-called the Casson–Lin invariant by applying the same idea to a certain *slice* (an algebraic subset) of the $\mathrm{SU}(2)$ -character variety of the knot exterior. Afterward, C. Herald generalized the Casson–Lin invariant by using gauge theoretic methods in [Her97]. M. Heusener and J. Kroll also have studied the same issue by topological methods in [HK98]. An invariant defined in the similar way to the Casson invariant is called the Casson type invariant. The Casson–Lin invariant can be regarded as a kind of the equivariant Casson invariant, which is defined by using the equivariant theory of n -fold cyclic branched covers Σ_n along a knot in Σ with covering transformations. It is known that this invariant is expressed by using the equivariant signature of the knot (for more information, refer to the result of O. Collin and N. Saveliev [CS01]).

Date: September 29, 2008.

2000 Mathematics Subject Classification. Primary: 57M27, 57M05, Secondary: 57M12.

Key words and phrases. Character varieties; branched covers; binary dihedral representations; knots; metabelian representations.

Now we can also consider the Casson type invariants for the $\mathrm{SL}_2(\mathbb{C})$ -character variety $X(M)$ of a manifold M , for example, the $\mathrm{SL}_2(\mathbb{C})$ -Casson invariant [Cur01]. However, the similar equivalent theory to Collin–Saveliev for $\mathrm{SL}_2(\mathbb{C})$ has not appear yet. So, in this article, we are interested in understanding the equivariant theory for the $\mathrm{SL}_2(\mathbb{C})$ -Casson invariant and thus understanding the Casson–Lin invariant for $\mathrm{SL}_2(\mathbb{C})$.

As a first step to understand these objects from the Casson–Lin invariant point of view, we would like to focus on *the slice* $S_0(\Sigma_K)$ at 0 of the $\mathrm{SL}_2(\mathbb{C})$ -character variety $X(\Sigma_K)$ of a knot exterior Σ_K for a knot in Σ , which corresponds to the slice used in the construction of the Casson–Lin invariant (refer to Definition 4.12 in Section 4.3 or the original article [Lin92]). We first find that *the metabelian representations* (refer to the very beginning of Section 4) form the fixed point set of an involution on the slice $S_0(\Sigma_K)$ (refer to Proposition 4.8 and Lemma 4.13). This gives a necessary and sufficient condition for irreducible representations to be metabelian. Then we can construct explicitly a map denoted by $\widehat{\Phi}_2$ from the slice $S_0(\Sigma_K)$ to the $\mathrm{SL}_2(\mathbb{C})$ -character variety $X(\Sigma_2)$ of a two-fold branched cover Σ_2 . By using $\widehat{\Phi}_2$, we can study the geometry of the slice $S_0(\Sigma_K)$ as follows. First, the map $\widehat{\Phi}_2$ turns out to have a property such as a covering map for a two-fold branched cover. If $\widehat{\Phi}_2$ is surjective, then the slice has a two-fold branched cover-like structure whose base space can be thought of as the $\mathrm{SL}_2(\mathbb{C})$ -character variety $X(\Sigma_2)$ and whose branched point set consists entirely of the characters of metabelian representations of the knot group (refer to Theorem 6.3, which is a main result in this article). We can observe that the map $\widehat{\Phi}_2$ is surjective for two-bridge knots and pretzel knots of type (p, q, r) (refer to Section 7).

In the above process, we find an explicit correspondence between the metabelian representations of a knot group $\pi_1(\Sigma_K)$ and the abelian ones for $\pi_1(\Sigma_2)$. In fact, there exists a map Φ_2 from the set of $\mathrm{SL}_2(\mathbb{C})$ -representations of $\pi_1(\Sigma_K)$ to that of $\pi_1(\Sigma_2)$ and the map Φ_2 gives the correspondence (see Proposition 6.2), where Φ_2 naturally induces $\widehat{\Phi}_2$. Then the characters of metabelian representations of the knot group coincides with the characters of *binary dihedral representations* of the knot group (refer to Proposition 4.8 and Corollary 4.10). This means that the representatives of conjugacy classes of metabelian representations can be taken by the binary dihedral representations. We demonstrate this for two-bridge knots (see Section 7). We also observe that all $\mathrm{SL}_2(\mathbb{C})$ -representations for a Seifert fibered manifold with three singular fibers are obtained by those of a pretzel knot group (see Section 7). The $\mathrm{SL}_2(\mathbb{C})$ -Casson invariant for Seifert fibered homology spheres have been calculated by surgeries in [BC06]. So, this is a starting point to understand the equivariant theory for the $\mathrm{SL}_2(\mathbb{C})$ -Casson invariant.

In fact, the above framework for a two-fold branched cover Σ_2 can be naturally extended to an n -fold cyclic branched cover Σ_n . Namely, we can extend Φ_2 to the map Φ_n from a certain set $R_{\xi(k/n)}(\Sigma_K)$ of $\mathrm{SL}_2(\mathbb{C})$ -representations of Σ_K to the set $R(\Sigma_n)$ of $\mathrm{SL}_2(\mathbb{C})$ -representations of $\pi_1(\Sigma_n)$ (see Section 5). Then we study relationships between their characters, i.e., the slice $S_{\xi(k/n)}(\Sigma_K)$ of $X(\Sigma_K)$ associated to $R_{\xi(k/n)}(\Sigma_K)$ and $X(\Sigma_n)$. In fact, such a relationship is already studied by Collin and Saveliev [CS01] in the case of the $\mathrm{SU}(2)$ -character varieties for cyclic branched covers under the assumption that a cyclic branched cover and its base manifold are both integral homology spheres. So we generalize their result to the case of $\mathrm{SL}_2(\mathbb{C})$ -character varieties for cyclic branched covers without any assumptions.

Our approach to this topic is rather more algebraic than Collin-Saveliev. The main tool is Lin's presentation, which is a special presentation of a knot group (refer to Section 3 or the original article [Lin01]). We see that Lin's presentation induces presentations of fundamental groups of covering spaces. By using this, we show that $\mathrm{SL}_2(\mathbb{C})$ -representations associated to $R_{\xi(k/n)}(\Sigma_K)$ induce $\mathrm{PSL}_2(\mathbb{C})$ -representations for the corresponding covering space Σ_n , where $\mathrm{PSL}_2(\mathbb{C}) = \mathrm{SL}_2(\mathbb{C})/\{\pm 1\}$. Lifting these $\mathrm{PSL}_2(\mathbb{C})$ -representations to $\mathrm{SL}_2(\mathbb{C})$, we can construct the desired map Φ_n (we also give a method to lift $\mathrm{PSL}_2(\mathbb{C})$ -representations to $\mathrm{SL}_2(\mathbb{C})$ -ones by using the abelianization homomorphism of a knot group). Then we investigate properties of the map Φ_n between representations. This framework extension is another main result in this article.

Organization. We explain the above statement in the following steps. In Section 2, we review some notions about representations and the characters and define some notations used throughout this article. Section 3 gives a review of Lin's presentation of a knot group and study its properties. In Section 4, we discuss the characters of metabelian representations, which tells that the metabelian representations are closely related to the binary dihedral representations. In Section 5, first we observe that what kinds of $\mathrm{SL}_2(\mathbb{C})$ -representations of a knot group induce $\mathrm{PSL}_2(\mathbb{C})$ -representations of covering spaces and next how to lift $\mathrm{PSL}_2(\mathbb{C})$ -representations to $\mathrm{SL}_2(\mathbb{C})$ -representations. In Section 6, we focus on the geometry of the slice $S_0(\Sigma_K)$ at 0, especially on the two-fold branched cover-like structure. In the final section, we show some applications and examples, which show that the map $\widehat{\Phi}_2$ is surjective for two-bridge knots and pretzel knots of type (p, q, r) .

2. PRELIMINARIES

2.1. General notations. We use the symbol Σ to denote an integral homology three-sphere and K to denote a knot in Σ . Then Σ_K stands for the knot exterior, i.e., it is obtained by removing an open tubular neighbourhood of K from Σ . Let us denote by Σ_n the n -fold cyclic branched cover of Σ along the branched set K and by C_n the n -fold cyclic cover of Σ_K . The following diagram expresses the relationship among these spaces,

$$\begin{array}{ccc} C_n & \longrightarrow & \Sigma_n \\ \downarrow & & \downarrow \\ \Sigma_K & \longrightarrow & \Sigma. \end{array}$$

The right arrows are inclusions and the down arrows are projections. When we consider several fundamental groups $\pi_1(\Sigma)$, $\pi_1(\Sigma_K)$, $\pi_1(C_n)$ and $\pi_1(\Sigma_n)$, we assume that the base points of Σ and Σ_K are given by the same point b which lives in the boundary torus of Σ_K and those of C_n and Σ_n are given by a lift b_n of b . We will abbreviate $\pi_1(\Sigma, b)$ to $\pi_1(\Sigma)$ and so on. We denote by μ a meridian of a knot K which is represented by a simple closed curve in the boundary torus.

2.2. Review on the $\mathrm{SL}_2(\mathbb{C})$ -character variety. We briefly review the $\mathrm{SL}_2(\mathbb{C})$ -representation space and the character variety. We consider $\mathrm{SL}_2(\mathbb{C})$ -representations for the fundamental group of a compact manifold M , i.e., homomorphisms from $\pi_1(M)$ into $\mathrm{SL}_2(\mathbb{C})$. We let

$$R(M) = \mathrm{Hom}(\pi_1(M); \mathrm{SL}_2(\mathbb{C}))$$

denote the space of $\mathrm{SL}_2(\mathbb{C})$ -representations for M . As usual, this space is endowed with the compact-open topology. Here $\pi_1(M)$ is assumed to have the discrete topology and the Lie group $\mathrm{SL}_2(\mathbb{C})$ is endowed with the usual one.

An representation $\rho : \pi_1(M) \rightarrow \mathrm{SL}_2(\mathbb{C})$ is called *abelian* if the image $\rho(\pi_1(M))$ is an abelian subgroup in $\mathrm{SL}_2(\mathbb{C})$. We say that ρ is *metabelian* if the commutator subgroup $[\pi_1(M), \pi_1(M)]$ is sent to an abelian subgroup in $\mathrm{SL}_2(\mathbb{C})$ by ρ . Of course, all abelian representations are metabelian.

It is well-known that there exist only the following two maximal abelian subgroups, up to conjugation, in $\mathrm{SL}_2(\mathbb{C})$:

$$\begin{aligned} \mathrm{Hyp} &:= \left\{ \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} \mid \lambda \in \mathbb{C}^* := \mathbb{C} \setminus \{0\} \right\}, \\ \mathrm{Para} &:= \left\{ \pm \begin{pmatrix} 1 & \omega \\ 0 & 1 \end{pmatrix} \mid \omega \in \mathbb{C} \right\}. \end{aligned}$$

Here Hyp means hyperbolic elements in $\mathrm{SL}_2(\mathbb{C})$ and Para means parabolic elements in $\mathrm{SL}_2(\mathbb{C})$. Hence if ρ is an abelian (resp. metabelian) $\mathrm{SL}_2(\mathbb{C})$ -representation, the image (resp. the image of $[\pi_1(M), \pi_1(M)]$) can be contained in Hyp or Para by taking conjugation.

An $\mathrm{SL}_2(\mathbb{C})$ -representation ρ is called *reducible* if there exists a *proper* subspace $U \subset \mathbb{C}^2$ such that $\rho(g)(U) \subset U$, for all $g \in \pi_1(M)$. Namely, there exists an element A of $\mathrm{SL}_2(\mathbb{C})$ such that $A\rho(g)A^{-1}$ is an upper triangular matrix for any $g \in \pi_1(M)$. Of course, any abelian representation is reducible (while the converse is false in general). Moreover by [CS83, Lemma 1.2.1], all reducible representations send the commutator subgroup $[\pi_1(M), \pi_1(M)]$ into the maximal abelian subgroup Para, up to conjugate. Hence reducible representations are metabelian. A representation is called *irreducible* if it is not reducible.

The group $\mathrm{SL}_2(\mathbb{C})$ acts on the representation space $R(M)$ by conjugation, however the naive quotient $R(M)/\mathrm{SL}_2(\mathbb{C})$ is not Hausdorff in general. Following [CS83], we will focus on the set of *characters* of $R(M)$. Associated to a representation $\rho \in R(M)$, its character $\chi_\rho : \pi_1(M) \rightarrow \mathbb{C}$ is defined by $\chi_\rho(g) = \mathrm{tr} \rho(g)$, where tr denotes the trace of matrices. It is well-known that $R(M)$ and the set of characters have the structure of complex algebraic affine varieties, see [CS83]. In this sense, the set of characters of $R(M)$ is called the character variety of M and denoted by $X(M)$. In some sense $X(M)$ is the “algebraic quotient” of $R(M)$ by the action by conjugation of $\mathrm{PSL}_2(\mathbb{C})$. It is well known that $R(M)$ and $X(M)$ have the structure of complex algebraic affine varieties, see [CS83].

Let $R^{\mathrm{irr}}(M)$ denote the subset of irreducible representations of $\pi_1(M)$, and $X^{\mathrm{irr}}(M)$ denote its image under the map $t : R(M) \rightarrow X(M)$, $t(\rho) = \chi_\rho$. Note that two irreducible representations of $\pi_1(M)$ in $\mathrm{SL}_2(\mathbb{C})$ with the same character are conjugate by an element of $\mathrm{SL}_2(\mathbb{C})$, see [CS83, Proposition 1.5.2].

Similarly, we write $X^{\mathrm{red}}(M)$ for the image of the set $R^{\mathrm{red}}(M)$ of reducible representations in $X(M)$. We also use $R^{\mathrm{ab}}(M)$ (resp. $R^{\mathrm{meta}}(M)$) for the set of abelian (resp. metabelian) representations and $X^{\mathrm{ab}}(M)$ (resp. $X^{\mathrm{meta}}(M)$) for the image by t . We will see that $X^{\mathrm{red}}(M) \subset X^{\mathrm{meta}}(M)$ and this inclusion is strict in general, i.e., there exist irreducible metabelian representations.

3. LIN'S PRESENTATION

We will consider character varieties of cyclic covers over a knot exterior. A knot group has several presentations, for example, Wirtinger presentation, braid presentation. X-S. Lin has introduced a presentation of a knot group by using a free Seifert surface and the induced Heegaard splitting of S^3 . We call this presentation *Lin's presentation*. We show that Lin's presentation induces presentations of the fundamental groups of covering spaces. Moreover we present nice properties of the induced presentations. These presentations are main tools for our investigation of character varieties in the rest of this article.

3.1. Lin's presentation for an integral homology sphere. Lin's presentation was given for a knot in S^3 , however, his construction can be extended for a knot in an integral homology three-sphere. To explain this extension, we review his construction.

First we review the definition of a free Seifert surface.

Definition 3.1. A Seifert surface S of a knot K is called free if $(N(S), \overline{\Sigma \setminus N(S)})$ is a Heegaard splitting of an integral homology three-sphere Σ . Here $N(S)$ is a closed tubular neighborhood of S in Σ .

Such a Seifert surface always exist in an integral homology sphere Σ as follows.

Lemma 3.2. *Let K be a knot in an integral homology three-sphere Σ . Then there exists a Heegaard splitting $\Sigma = H_1 \cup_F H_2$ such that $H_1 = S \times [-1, 1]$ and $K = \partial S$ is a separating curve on F .*

For more information about the existence of a free Seifert surface and the induced Heegaard decomposition of an integral homology three-sphere, for example, see [Sav99, Lecture 17].

A presentation of $\pi_1(\Sigma_K)$ is obtained from the Heegaard splitting associated to a free Seifert surface.

Lemma 3.3 (Lemma 2.1 in [Lin01]). *Assume that S is a free Seifert surface for K in Σ . Let $\Sigma = H_1 \cup H_2$ be the Heegaard splitting associated to S . The fundamental group $\pi_1(H_2)$ is a free group and we denote by $x_1, \dots, x_{2g}$ a basis of $\pi_1(H_2)$.*

Then we have the following presentation of $\pi_1(\Sigma_K)$:

$$(1) \quad \pi_1(\Sigma_K) = \langle x_1, \dots, x_{2g}, \mu \mid \mu \alpha_i \mu^{-1} = \beta_i, i = 1, \dots, 2g \rangle,$$

where α_i, β_i are words in $x_1, \dots, x_{2g}$, determined by the spine of S , and g is the genus of S .

We call the presentation (1) Lin's presentation.

Remark 3.4. Note that every x_i is an element in the commutator subgroup of $\pi_1(\Sigma_K)$.

This can be shown by van Kampen's theorem as follows. As in Lemme 3.2, the integral homology sphere Σ is decomposed into two handle bodies H_1 and H_2 where $H_1 = S \times [-1, 1]$. To make a decomposition of Σ_K from this Heegaard splitting such that the intersection of two parts is connected, we push K into H_1 slightly and remove a tubular neighbourhood of the perturbed knot from H_1 . To use van Kampen's theorem, we describe this decomposition more precisely. Let W_{2g} be a spine of S i.e., a bouquet with a base point of circles in S which is a deformation

retract of S , where g is the genus of S . We denote by $a_1, a_2, \dots, a_{2g-1}, a_{2g}$ the oriented circles in W_{2g} such that they form a symplectic basis of S with that order. The base point of W_{2g} is denoted by $*$. Let K' be a simple closed curve on $S \times \{0\}$ parallel to $\partial S = K$ and disjoint from W_{2g} . Identifying the knots K' and K , we can consider the decomposition $\Sigma_K = (H_1 \setminus \text{int } N(K')) \cup H_2$. The piece $H_1 \setminus \text{int } N(K')$ is homotopy equivalent to $T^2 \cup S$ where $T^2 = \partial N(K')$ and the intersection $T^2 \cap S$ is the longitude. Hence the fundamental group $\pi_1(H_1 \setminus \text{int } N(K'))$ is generated by the homotopy classes of $a_1, \dots, a_{2g}$ and the meridian.

By van Kampen's theorem, the knot group $\pi_1(\Sigma_K)$ is isomorphic to the amalgamated product of $\pi_1(H_1 \setminus \text{int } N(K'))$ and $\pi_1(H_2)$. Hence the knot group is generated by $x_1, \dots, x_{2g}$, the homotopy classes of $a_1, \dots, a_{2g}$ and the meridian. Then the relations are given by elements of $\pi_1(\partial H_1)$. We can simplify the presentation of this amalgamated product. Let $H_1 = S \times [-1, 1]$ be a bicollar of S given by the positive normal direction of S such that $S = S \times \{0\}$. For $i = 1, \dots, 2g$, let $a_i^\pm = a_i \times \{\pm 1\}$ be circles in $W_{2g} \times \{\pm 1\}$ respectively. There exists an arc c on ∂H_1 , uniquely determined up to isotopy, connecting the two point $\{*\} \times \{\pm 1\}$ and its interior has no intersection with $W_{2g} \times \{\pm 1\}$. The circle $c \cup (\{*\} \times [-1, 1])$ can be thought of as a meridian μ of K' .

Now, the homotopy class of a_i^+ (resp. a_i^-) is expressed as a word in $x_1, \dots, x_{2g}$ for each i . We denote by α_i (resp. β_i) the word for a_i^+ (resp. a_i^-). Since each simple closed curve a_i^+ belongs to the same homotopy class of a_i , the knot group $\pi_1(\Sigma_K)$ is generated by $x_1, \dots, x_{2g}, \mu$. Since $x_1, \dots, x_{2g}$ are generators of a free group, the relations of $\pi_1(\Sigma_K)$ are given by disks in $H_1 \setminus N(K')$. We assume that such a disk D^2 meets $S \times \{0\}$ transversely and the boundary is pushed toward $\partial(H_1 \setminus \text{int } N(K'))$. The homotopy class of the intersection D^2 and $S \times \{0\}$ is give by a word of $a_1, \dots, a_{2g}$. This D^2 defines the relation by replacing each a_i with $\mu \alpha_i \mu^{-1} \beta_i^{-1}$ in the word of $a_1, \dots, a_{2g}$. Since the relations are generated by $\mu \alpha_i \mu^{-1} \beta_i^{-1}$, we obtain the presentation in Lemma 3.3.




FIGURE 1. Seifert surface

Let us denote by $v_{i,j}$ the exponent sum of x_j in α_i and $u_{i,j}$ the exponent sum of x_i in β_i . We set two $2g \times 2g$ matrices $V := (v_{i,j})$ and $U := (u_{i,j})$ with integer entries.

For a knot in S^3 , Lin has introduced special free Seifert surface, called *regular* (see [Lin01]). A free Seifert surface S is called regular if it has a spine where embedding in S^3 induced by S is isotopic to the standard embedding. By setting appropriate orientations on a_i and x_j , we obtain the following standard properties of V and U :

- (1) U is so-called the Seifert matrix $Q = (Q_{i,j})$, $Q_{i,j} = lk(a_i, a_j^+)$, for S ;

(2) $V = {}^tU$, where tU is the transpose matrix of V .

Note that in Lin's paper the matrix $(lk(a_i^+, a_j))_{i,j}$ is used as the Seifert matrix, which is different from ours $(lk(a_i, a_j^+))_{i,j}$. This gives rise to differences between Lin's convention of matrices U and V and ours.

Unfortunately, regular Seifert surfaces are defined only for knots in S^3 . However, V and U are expressed by using the Seifert matrix for a general free Seifert surface of a knot in an integral homology sphere Σ . To describe V and U for a free Seifert surface in Σ , we need one more $2g \times 2g$ matrix T which is obtained from the Morse complex $C_*^{Morse}(\Sigma; \mathbb{Z})$ associated to the Heegaard splitting as follows.

Let S be a free Seifert surface for a knot K in Σ . Set $\Sigma = H_1 \cup H_2$ as a Heegaard splitting associated to S . The set of cores of 1-handle in H_1 are given by $\{a_1, \dots, a_{2g}\}$ in the spine W_{2g} . We denote by D_i a meridian disk for a_i . The set of cores of 2-handles forms that of generators $\{x_1, \dots, x_{2g}\}$ in $\pi_1(H_2)$. We denote by D'_i a meridian disk for x_i .

It is known that there exists a Morse function f compatible with the Heegaard splitting (see for instance [OS04]). Here "compatible" means that $f : Y \rightarrow [0, 3]$ is a self-indexing Morse function on Σ with one minimum and one maximum and f induces the Heegaard decomposition with surface $S = f^{-1}(3/2)$, $H_1 = f^{-1}([0, 3/2])$, $H_2 = f^{-1}([3/2, 3])$. The attaching circles ∂D_i and $\partial D'_i$ are the intersections of S with the ascending and descending manifolds for the index one and two critical points respectively.

The intersection point between the meridian disk D_i and the core a_i gives a critical point with index one. We denote by q_i this critical point. The intersection D'_i with x_i gives a critical point with index two. We denote it by p_i . Each 2-handle are attached to H_1 along the image $\partial D'_i$ in ∂H_1 . We set an integer $t_{k,j}$ as the intersection number ∂D_j with $\partial D'_k$ in ∂H_1 . The matrix $(t_{k,j})$ gives the boundary operator from $C_2^{Morse}(\Sigma; \mathbb{Z})$ to $C_1^{Morse}(\Sigma; \mathbb{Z})$. The correspondence between the bases $(p_1, \dots, p_{2g})$ and $(q_1, \dots, q_{2g})$ is expressed as

$$\partial(p_1, \dots, p_{2g}) = (q_1, \dots, q_{2g})T,$$

where T is the $2g \times 2g$ -matrix $(t_{k,j})$.




FIGURE 2. Heegaard splitting

Remark 3.5. Since Σ is an integral homology three-sphere, the representation matrix T is invertible. In particular the determinant $\det(T)$ is equal to ± 1 .

Proposition 3.6. *We suppose that S is a free Seifert surface of a knot K . Let V and U be the matrices defined by the relations α_i and β_i in Lin's presentation (1) of the knot group. Then the matrices V and U are expressed as*

$$V = {}^tQT, \quad U = QT,$$

where Q is the Seifert matrix for S and T is the matrix corresponding the boundary operator from $C_2^{Morse}(\Sigma; \mathbb{Z})$ to $C_1^{Morse}(\Sigma; \mathbb{Z})$.

Proof. From the definition, the integer $v_{i,j}$ is given by the intersection number a_i^+ with $\partial D'_j$ in ∂H_1 . Since $\partial D'_j$ is homologue to $\sum_{k=1}^{2g} t_{k,j} a_k$, by using linking number, the entry $v_{i,j}$ in V is expressed as

$$(2) \quad v_{i,j} = lk(a_i^+, \sum_{k=1}^{2g} t_{k,j} a_k).$$

Here $t_{k,j}$ is the entries in the matrix T . The right hand side of (2) turns into

$$\begin{aligned} v_{i,j} &= lk(a_i^+, \sum_{k=1}^{2g} t_{k,j} a_k) \\ &= \sum_{k=1}^{2g} t_{k,j} lk(a_i^+, a_k) \\ &= \sum_{k=1}^{2g} t_{k,j} Q_{k,i}. \end{aligned}$$

This means that $V = {}^tQT$. Similarly we can prove that $U = QT$. □ □

Remark 3.7. The Alexander polynomial $\Delta_K(t)$ is given by $\det(tU - V)$.

Remark 3.8. If K is a knot in S^3 and S is regular, then the matrix T is the identity matrix.

3.2. Presentation of $\pi_1(C_n)$. The n -fold cyclic cover C_n over Σ_K has the covering transformations. The covering transformations form a cyclic group with order n . This action induces the automorphism τ_n of $\pi_1(C_n)$,

$$\begin{aligned} \tau_n : \pi_1(C_n, b_n) &\rightarrow \pi_1(C_n, b_n), \\ [l] &\mapsto [\tilde{m} \cdot \tau(l) \cdot \tilde{m}^{-1}], \end{aligned}$$

where b_n is a lift of the base point b of $\pi_1(\Sigma_K)$, l denotes a closed loop in C_n , τ is the covering transformation and $\tilde{m}$ is a lift of the meridian μ with the initial point b_n . Let $p_n : \pi_1(C_n, b_n) \rightarrow \pi_1(\Sigma_K, b)$ be the homomorphism induced by the projection from C_n to Σ_K . Then we denote by μ_n a meridian of C_n such that $p_n(\mu_n) = \mu^n$. Note that the covering transformation does not preserve base point b_n .

Remark 3.9. It follows from the definition of τ_n that

- (1) for every element γ in $\pi_1(C_n)$, the action of τ_n^n sends γ to $\mu_n \gamma \mu_n^{-1}$ where τ_n^k denotes k -times composition of τ_n ; and
- (2) the homomorphism τ_n sends μ_n to itself in $\pi_1(C_n)$.

Lin's presentation is useful to describe the fundamental groups of the cyclic cover C_n over Σ_K . Indeed, we can construct C_n by cutting Σ_K along a Seifert surface and gluing n copies. Then every closed loop on the Seifert surface can be lifted to C_n .

Lemma 3.10. *We suppose that the following Lin's presentation of $\pi_1(\Sigma_K)$ is given.*

$$\pi_1(\Sigma_K) = \langle x_1, \dots, x_{2g}, \mu \mid \mu \alpha_i \mu^{-1} = \beta_i \rangle.$$

Then $\pi_1(C_n)$ is presented as

$$(3) \quad \left\langle \begin{array}{l} \tilde{x}_1, \dots, \tilde{x}_{2g}, \tau_n(\tilde{x}_1), \dots, \tau_n(\tilde{x}_{2g}), \\ \dots, \tau_n^{n-1}(\tilde{x}_1), \dots, \tau_n^{n-1}(\tilde{x}_{2g}), \mu_n \end{array} \mid \begin{array}{l} \mu_n \tilde{\alpha}_i^{(0)} \mu_n^{-1} = \tilde{\beta}_i^{(n-1)}, \tilde{\alpha}_i^{(j)} = \tilde{\beta}_i^{(j-1)} \\ (1 \leq i \leq 2g, 1 \leq j \leq n-1) \end{array} \right\rangle,$$

where $\tilde{x}_i$ is a lift of x_i to C_n , $\tilde{\alpha}_i^{(j)}$ and $\tilde{\beta}_i^{(j)}$ are the words by replacing x_k in α_i and β_i with $\tau_n^j(\tilde{x}_k)$.

Proof. The presentation (3) follows from the construction of C_n by cutting Σ_K along the free Seifert surface S and gluing copies along the surfaces. First we separate Σ into two handle bodies $H_1 = S \times [-1, 1]$ and H_2 as in Lemma 3.2 and make a simple closed curve K' on $S \times \{0\}$ parallel to K by pushing K slightly into H_1 . We denote by S' a compact surface obtained by shrinking S so that the boundary coincides with K' . Cutting H_1 along S' and gluing n -copies, we have the cyclic cover $\tilde{H}_1$ of H_1 . Since the cyclic cover $\tilde{H}_1$ is deformed to the union T^2 and n -copies of S' , the fundamental group $\pi_1(\tilde{H}_1)$ is generated by the homotopy classes of all lifts of a_i ($1 \leq i \leq 2g$) and μ_n where $a_1, \dots, a_{2g}$ are circles in a spine of S' . Next gluing n -copies of H_2 to $\tilde{H}_1$ as in Figure 3, we obtain the n -fold cyclic cover of Σ_K and the presentation (3) by van Kampen's theorem. $\square$ $\square$




FIGURE 3. Decomposition of C_n

The presentation of $\pi_1(C_n)$ in Lemma 3.10 induces a presentation of $H_1(C_n; \mathbb{Z})$ as follows. It is known that this presentation of $H_1(C_n; \mathbb{Z})$ can be also proved by using Mayer-Vietoris argument and the Alexander duality, see [Lic97, Chapter 9].

Lemma 3.11. *We keep the notations in Proposition 3.6 and Lemma 3.10. Let Z be a free abelian group generated by $2ng$ -elements as follows:*

$$Z := \left(\oplus_{i=1}^{2g} \mathbb{Z}[\tilde{x}_i] \right) \oplus \left(\oplus_{i=1}^{2g} \mathbb{Z}[\tau_n(\tilde{x}_i)] \right) \cdots \oplus \left(\oplus_{i=1}^{2g} \mathbb{Z}[\tau_n^{n-1}(\tilde{x}_i)] \right).$$

Then the presentation of $H_1(C_2; \mathbb{Z})$ is given by

$$Z \xrightarrow{\begin{pmatrix} A \\ \mathbf{0} \end{pmatrix}} Z \oplus \mathbb{Z}[\mu_n] \rightarrow H_1(C_n; \mathbb{Z}),$$

9

where

$$A = \begin{pmatrix} V & -U & \cdots & 0 & 0 \\ 0 & V & -U & \cdots & 0 \\ 0 & 0 & V & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ -U & \cdots & 0 & 0 & V \end{pmatrix}.$$

Moreover if we set $\mathbf{x}_j = {}^t([\tau_n^j(\tilde{x}_1)], \dots, [\tau_n^j(\tilde{x}_{2g})])$, then we have the following relation in $H_1(C_n; \mathbb{Z})$:

$$\mathbf{x}_0 + \cdots + \mathbf{x}_{n-1} = \mathbf{0}.$$

Proof. The above presentation is given by the abelianization of the presentation in Lemma 3.10. So we omit the details. Here we show the relation $\mathbf{x}_0 + \cdots + \mathbf{x}_{n-1} = \mathbf{0}$. From the presentation of $H_1(C_n; \mathbb{Z})$, we have the following relation:

$$\begin{cases} V\mathbf{x}_0 & -U\mathbf{x}_{n-1} & = & \mathbf{0} \\ -U\mathbf{x}_0 & +V\mathbf{x}_1 & = & \mathbf{0} \\ & \vdots & & \vdots \\ & -U\mathbf{x}_{n-2} & +V\mathbf{x}_{n-1} & = & \mathbf{0}. \end{cases}$$

By taking sum on the both side, we have

$$(V - U)(\mathbf{x}_0 + \cdots + \mathbf{x}_{n-1}) = \mathbf{0}.$$

By Proposition 3.6 and the property of the Alexander polynomial, the absolute value $|\det(V - U)|$ is equal to $|\Delta_K(1)| = 1$. Hence we have the equation that $\mathbf{x}_0 + \cdots + \mathbf{x}_{n-1} = \mathbf{0}$. $\square$ $\square$

3.3. Presentation of $\pi_1(\Sigma_n)$. We denote by π_n the induced homomorphism from $\pi_1(C_n)$ to $\pi_1(\Sigma_n)$ induced by the inclusion from C_n onto Σ_n . Now we have the following diagrams on the knot exterior and its covering spaces:

$$\begin{array}{ccc} C_n & \longrightarrow & \Sigma_n \\ \downarrow & & \downarrow \\ \Sigma_K & \longrightarrow & \Sigma, \end{array} \quad \begin{array}{ccc} \pi_1(C_n) & \xrightarrow{\pi_n} & \pi_1(\Sigma_n) \\ p_n \downarrow & & \downarrow \\ \pi_1(\Sigma_K) & \xrightarrow{\pi} & \pi_1(\Sigma). \end{array}$$

Since Σ_n is obtained from C_n by gluing $D^2 \times S^1$ and the universality of the amalgamated product, we have the following diagram:

$$(4) \quad \begin{array}{ccccc} & & \pi_1(D^2 \times S^1) & \searrow & \\ \pi_1(T^2) & \nearrow & & & \\ & & \pi_1(C_n) & \nearrow & \\ & & & & \pi_1(D^2 \times S^1) *_{\pi_1(T^2)} \pi_1(C_n) \xrightarrow{f} \pi_1(\Sigma_n). \end{array}$$

By van Kampen's theorem, the above f is isomorphism. The following Lemma is a fundamental fact in the combinatorial group theory.

Lemma 3.12. *Suppose that φ in the following diagram is surjective:*

$$\begin{array}{ccccc} & & G & \xrightarrow{i_G} & \\ K & \xrightarrow{\varphi} & & & G *_K H. \\ & \searrow \psi & H & \xrightarrow{i_H} & \end{array}$$

Then i_H is also surjective and the kernel of i_H is given by $|\psi(\ker \varphi)|$ where $|T|$ is the normal closure of a subset T . In particular, we have the isomorphism $H/|\psi(\ker \varphi)| \simeq G *_K H$.

Since the homomorphism $\pi_1(T^2) \rightarrow \pi_1(D^2 \times S^1)$ is surjective in the diagram (4) and the kernel is generated by a meridian, we have the isomorphism:

$$(5) \quad \pi_1(C_n)/|\langle \mu_n \rangle| \simeq \pi_1(\Sigma_n).$$

where μ_n is the meridian such that $p_n(\mu_n) = \mu^n$. Hence we can regard the homomorphism π_n as the projection from $\pi_1(C_n)$ onto $\pi_1(C_n)/|\langle \mu_n \rangle|$. Similarly, we have the following isomorphism for $\pi_1(\Sigma)$:

$$(6) \quad \pi_1(\Sigma_K)/|\langle \mu \rangle| \simeq \pi_1(\Sigma).$$

We review one lemma about the normal closure.

Lemma 3.13. *Let G be a group. For a subset T in G , the normal closure $|T|$ is expressed as*

$$|T| = \{(g_1^{-1} t_1^{\epsilon_1} g_1) \cdots (g_k^{-1} t_k^{\epsilon_k} g_k) \mid k \geq 0, g_i \in G, t_i \in T, \epsilon_i = \pm 1 (1 \leq i \leq k)\}.$$

Corollary 3.14. *If a homomorphism $\psi : G \rightarrow H$ satisfies that $\psi(T) = 1_H$ where 1_H is the identity element of H , then ψ induces a homomorphism from $G/|T|$ to H .*

We have presentations of $\pi_1(C_n)$ and $H_1(C_n; \mathbb{Z})$ in Lemma 3.10 and 3.11. From the above relation between $\pi_1(C_n)$ and $\pi_1(\Sigma_n)$, we obtain the following presentations of $\pi_1(\Sigma_n)$ and $H_1(\Sigma_n; \mathbb{Z})$.

Lemma 3.15. *For a Lin's presentation of $\pi_1(\Sigma_K)$,*

$$\pi_1(\Sigma_K) = \langle x_1, \dots, x_{2g}, \mu \mid \mu \alpha_i \mu^{-1} = \beta_i \rangle,$$

$\pi_1(\Sigma_n)$ is presented as

$$(7) \quad \left\langle \begin{array}{l} \tilde{x}_1, \dots, \tilde{x}_{2g}, \tau_n(\tilde{x}_1), \dots, \tau_n(\tilde{x}_{2g}), \\ \dots, \tau_n^{n-1}(\tilde{x}_1), \dots, \tau_n^{n-1}(\tilde{x}_{2g}) \end{array} \mid \begin{array}{l} \tilde{\alpha}_i^{(0)} = \tilde{\beta}_i^{(n-1)}, \tilde{\alpha}_i^{(j)} = \tilde{\beta}_i^{(j-1)} \\ (1 \leq i \leq 2g, 1 \leq j \leq n-1) \end{array} \right\rangle,$$

where $\tilde{x}_i$ is a lift of x_i to C_n , $\tilde{\alpha}_i^{(j)}$ (resp. $\tilde{\beta}_i^{(j)}$) is the words by replacing x_k in α_i (resp. β_i) with $\tau_n^j(\tilde{x}_k)$. The homology group $H_1(\Sigma_n; \mathbb{Z})$ is expressed as

$$Z \xrightarrow{A} Z \rightarrow H_1(\Sigma_n; \mathbb{Z}),$$

where Z and A are as in Lemme 3.11. Moreover for $\mathbf{x}_j = {}^t([\tau_n^j(\tilde{x}_1)], \dots, [\tau_n^j(\tilde{x}_{2g})])$, the following relation holds in $H_1(C_n; \mathbb{Z})$:

$$\mathbf{x}_0 + \dots + \mathbf{x}_{n-1} = \mathbf{0}.$$

Remark 3.16. The group $\pi_1(\Sigma_n)$ is obtained by adding a relation $\mu_n = 1$. Then $\rho \in R(C_n)$ can be considered as an element $\rho \in R(\Sigma_n)$ if and only if ρ satisfies that $\rho(\mu_n) = \mathbf{1}$.

4. ON THE CHARACTERS OF METABELIAN REPRESENTATIONS

We consider metabelian representations of $\pi_1(\Sigma_K)$ in this section. A metabelian representation is a homomorphism from a group $\pi_1(\Sigma_K)$ into $\mathrm{SL}_2(\mathbb{C})$ such that it sends $[\pi_1(\Sigma_K), \pi_1(\Sigma_K)]$ to an abelian subgroup in $\mathrm{SL}_2(\mathbb{C})$.

4.1. Reducible representations. From the definition, every reducible representation is conjugate to a representation whose image consists of upper triangular matrices in $\mathrm{SL}_2(\mathbb{C})$.

Remark 4.1. If A and B are upper triangular $\mathrm{SL}_2(\mathbb{C})$ -elements, then the commutator $[A, B]$ turns into a parabolic element. Hence we can see that the commutator subgroup of the image of any reducible representation is an abelian subgroup in $\mathrm{SL}_2(\mathbb{C})$. All reducible representations are metabelian. The set $R^{\mathrm{meta}}(\Sigma_K)$ of metabelian representations is decompose to the union

$$\{\rho \in R^{\mathrm{irr}}(\Sigma_K) \mid \rho : \text{metabelian}\} \cup R^{\mathrm{red}}(\Sigma_K).$$

Furthermore we have two kinds of reducible representations, one is abelian and another is non-abelian. Every abelian representation factors through the abelianization $H_1(\Sigma_K; \mathbb{Z})$. Since the homology group $H_1(\Sigma_K; \mathbb{Z})$ is generated by the homology class of the meridian μ , each abelian representation is determined by the image of the meridian μ . For a given $\lambda \in \mathbb{C}^*$, we denote by φ_λ the abelian representation given by

$$\varphi_\lambda : \pi_1(\Sigma_K) \ni \mu \mapsto \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C}).$$

By taking conjugation, we can assume that the image of any reducible representation consists of upper triangular $\mathrm{SL}_2(\mathbb{C})$ -elements. This means that for every reducible representation there exists a complex number λ such that the character of φ_λ is the same as that of the reducible one. For non-abelian reducible representations, it is known that the following fact holds.

Lemma 4.2 ([HPS01]). *There exists a reducible non-abelian representation*

$$\rho_\lambda : \pi_1(\Sigma_K) \rightarrow \mathrm{SL}_2(\mathbb{C})$$

such that $\chi_{\rho_\lambda} = \chi_{\varphi_\lambda}$ if and only if $\Delta_K(\lambda^2) = 0$.

This is a well-known result of Burde [Bur67] and de Rham [dR67] if Σ_K is the knot exterior of a knot in S^3 .

4.2. The characters of metabelian and binary dihedral representations. In this subsection, we focus on irreducible metabelian representation of a knot group. First we review binary dihedral representations of a knot group $\pi_1(\Sigma_K)$. As we will see, binary dihedral representations form a subset of metabelian representations and give representatives of conjugacy classes for irreducible metabelian $\mathrm{SL}_2(\mathbb{C})$ -representations.

The Lie group $\mathrm{SU}(2)$ is a subgroup in $\mathrm{SL}_2(\mathbb{C})$ and we can identify $\mathrm{SU}(2)$ with the set of unit quaternions as follows:

$$\mathrm{SU}(2) = \{a\mathbf{1} + b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \mid a, b, c, d \in \mathbb{R}, a^2 + b^2 + c^2 + d^2 = 1\},$$

where

$$\mathbf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \mathbf{i} = \begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix}, \mathbf{j} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \mathbf{k} = \begin{pmatrix} 0 & \sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix}.$$

The binary dihedral subgroup N in $\mathrm{SU}(2)$ is defined as $N = S_A^1 \cup S_B^1$ where $S_A^1 := \{a + b\mathbf{i} \mid a, b \in \mathbb{R}, a^2 + b^2 = 1\}$ and $S_B^1 := \{c\mathbf{j} + d\mathbf{k} \mid c, d \in \mathbb{R}, c^2 + d^2 = 1\}$. A binary dihedral representation is a homomorphism from $\pi_1(\Sigma_K)$ into N .

Let g, h be elements in N . The products gh and hg have the following property:

- if $g, h \in S_A^1$, then $gh = hg \in S_A^1$;
- if $g \in S_A^1$ and $h \in S_B^1$, then $gh, hg \in S_B^1$; and
- if $g, h \in S_B^1$, then $gh, hg \in S_B^1$.

If we consider a knot K in S^3 , then for a Wirtinger presentation of $\pi_1(E_K)$,

$$\pi_1(E_K) = \langle y_1, \dots, y_k \mid y_{j_l}^{\epsilon_l} y_l y_{j_l}^{-\epsilon_l} = y_{l+1} \ (l = 1, \dots, k-1), \epsilon_l \in \{\pm 1\} \rangle,$$

every irreducible binary dihedral representation sends each generator y_l to the following $SU(2)$ -element:

$$e^{i\theta_l} \mathbf{j} = \cos \theta_l \cdot \mathbf{j} + \sin \theta_l \cdot \mathbf{k},$$

where $e^{i\theta} = \cos \theta \cdot \mathbf{1} + \sin \theta \cdot \mathbf{i}$. Note that θ_l satisfies $2\theta_{j_l} = \theta_l + \theta_{l+1}$ by the relation in the Wirtinger presentation.

All binary dihedral representations are metabelian. This is shown as follows.

Proposition 4.3. *The commutator subgroup $[N, N]$ of N is an abelian subgroup in $SU(2)$. In particular, all binary dihedral representations are metabelian.*

Proof. Every element in N can be expressed as the form $e^{i\theta} \mathbf{j}^\delta$ where $\delta \in \{0, \pm 1\}$. By direct calculations, it follows that $e^{i\theta} \mathbf{j} = \mathbf{j} e^{-i\theta}$. For any g_1 and g_2 in N , we put $g_l = e^{i\theta_l} \mathbf{j}^{\delta_l}$, where $\delta_l \in \{0, \pm 1\}$ ($l = 1, 2$). Then the sum of exponents of $\mathbf{j}$ in $[g_1, g_2]$ is zero. Hence the commutator $[g_1, g_2]$ can be expressed as the form $e^{i\theta}$. This shows that the commutator subgroup $[N, N]$ is contained in the maximal abelian subgroup Hyp. $\square$ $\square$

Though binary dihedral representations are contained in metabelian representations, the characters of binary dihedral representations gives all those of metabelian ones.

Proposition 4.4. *Every irreducible metabelian $SL_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_K)$ is conjugate to an irreducible binary dihedral representation. Hence we have the following equation:*

$$\{\chi_\rho \mid \rho : \text{irreducible metabelian}\} = \{\chi_\rho \mid \rho : \text{irreducible binary dihedral}\}.$$

Moreover the number of conjugacy classes of irreducible metabelian representations is given by

$$\frac{|\Delta_K(-1)| - 1}{2},$$

where $\Delta_K(t)$ is the Alexander polynomial of K .

For a knot in S^3 , we can find this relation in the proof of [Nag07, Proposition 1.1 and Theorem 1.2]. The proof is given by using Lin's presentation. We also use Lin's presentation for a knot in an integral homology sphere Σ to prove Proposition 4.4.

Proof of Proposition 4.4. We suppose that an $SL_2(\mathbb{C})$ -representation ρ is irreducible and metabelian. Then it follows from [Nag07] that for the generators in Lin's presentation (1), by taking conjugation, we have

$$\rho(x_i) = \begin{pmatrix} \alpha_i & 0 \\ 0 & \alpha_i^{-1} \end{pmatrix}, \quad \rho(\mu) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

If we set the matrix $(w_{i,j}) = V + U$, as in the proof of [Nag07, Proposition 1.1 and Theorem 1.2], then the relations in Lin's presentation give the following equations:

$$\alpha_1^{w_{i,1}} \cdots \alpha_{2g}^{w_{i,2g}} = 1, \text{ for } i = 1, \dots, 2g.$$

Writing $\alpha_i = r_i e^{\sqrt{-1}\theta_i}$, we have

$$(w_{i,j}) \begin{pmatrix} \log r_1 \\ \vdots \\ \log r_{2g} \end{pmatrix} = \mathbf{0}, \quad (w_{i,j}) \begin{pmatrix} \theta_1 \\ \vdots \\ \theta_{2g} \end{pmatrix} \equiv \mathbf{0} \pmod{2\pi\mathbb{Z}}.$$

By Proposition 3.6 and Remark 3.7, the determinant $\det(V+U)$ is equal to $\Delta_K(-1)$. Hence all r_i are 1 and $(\theta_1, \dots, \theta_{2g})$ has $|\Delta_K(-1)|$ -solutions modulo $2\pi\mathbb{Z}$.

The solutions except the trivial one $(\alpha_1, \dots, \alpha_{2g}) = (1, \dots, 1)$ give irreducible representations. Conjugations preserving the form (9) just interchange the diagonal components in $\rho(x_i)$. The number of conjugacy classes is given by $|\Delta_K(-1) - 1|/2$. $\square$

On the other hand, every character of reducible representations is given by that of abelian one whose image is contained in Hyp. Moreover all abelian representations whose image is contained in Hyp are binary dihedral. Summarizing, we have the next corollary.

Corollary 4.5. *Every characters of an $\mathrm{SL}_2(\mathbb{C})$ -metabelian representation is given by that of a binary dihedral one.*

4.3. An involution on the $\mathrm{SL}_2(\mathbb{C})$ -character variety. Next we investigate the characters of irreducible metabelian $\mathrm{SL}_2(\mathbb{C})$ -representations via an action on the character variety. Let α be the abelianization homomorphism of $\pi_1(\Sigma_K)$ such that

$$\begin{aligned} \alpha : \pi_1(\Sigma_K) &\rightarrow \mathbb{Z} = \langle t \rangle, \\ \mu &\mapsto t, \end{aligned}$$

where $\mathbb{Z} = \langle t \rangle$ is a multiplicative group generated by one variable t . Here we are using the identification $\pi_1(\Sigma_K)/[\pi_1(\Sigma_K), \pi_1(\Sigma_K)] \simeq H_1(\Sigma_K; \mathbb{Z}) \simeq \mathbb{Z}$.

Definition 4.6. We define a map ι on the $\mathrm{SL}_2(\mathbb{C})$ -representation space $R(\Sigma_K)$ as follows:

$$\begin{aligned} \iota : R(\Sigma_K) &\rightarrow R(\Sigma_K), \\ \rho &\mapsto \alpha_{-1} \cdot \rho, \end{aligned}$$

where α_{-1} is the following composition of α and an evaluation map from $\mathbb{Z}$ into $\{\pm 1\}$:

$$\begin{array}{ccc} \pi_1(\Sigma_K) & \xrightarrow{\alpha} & \mathbb{Z} = \langle t \rangle \\ & \searrow \alpha_{-1} & \downarrow t=-1 \\ & & \{\pm 1\}. \end{array}$$

This map induces the following map $\hat{\iota}$ on the $\mathrm{SL}_2(\mathbb{C})$ -character variety $X(\Sigma_K)$:

$$\begin{aligned} \hat{\iota} : X(\Sigma_K) &\rightarrow X(\Sigma_K), \\ \chi_\rho &\mapsto \chi_{\iota(\rho)}. \end{aligned}$$

Remark 4.7. By definition, both ι and $\hat{\iota}$ are involutions.

The involution $\hat{\iota}$ has the following fixed point set.

Proposition 4.8. *The fixed point set of $\hat{\iota}$ in $X^{irr}(\Sigma_K)$ is the set of characters of irreducible metabelian $\mathrm{SL}_2(\mathbb{C})$ -representations, i.e.,*

$$X^{irr}(\Sigma_K)^{\hat{\iota}} = \{\chi_\rho \mid \rho : \text{irreducible metabelian}\}$$

We need the following elementary lemma to prove Proposition 4.8.

Lemma 4.9. *We set A and C as*

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad C = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

If the trace of AC is zero, then $b = c$.

Poof of Proposition 4.8. We use a Lin's presentation for the knot group:

$$(8) \quad \pi_1(\Sigma_K) = \langle x_1, \dots, x_{2g}, \mu \mid \mu \alpha_i \mu^{-1} = \beta_i \ (i = 1, \dots, 2g) \rangle.$$

First we show that for every irreducible metabelian $\mathrm{SL}_2(\mathbb{C})$ -representation ρ the image $\iota(\rho)$ is conjugate to ρ .

It follows from Proposition 4.4 that after taking suitable conjugation, the generators in the presentation (8) are sent to the following matrices by ρ :

$$(9) \quad \rho(x_i) = \begin{pmatrix} \lambda_i & 0 \\ 0 & \lambda_i^{-1} \end{pmatrix}, \quad \rho(\mu) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \mathbf{j}.$$

Since every x_i is contained in the commutator subgroup of $\pi_1(\Sigma_K)$, the images of these generators by $\iota(\rho)$ are given by

$$\iota(\rho)(x_i) = \begin{pmatrix} \lambda_i & 0 \\ 0 & \lambda_i^{-1} \end{pmatrix}, \quad \iota(\rho)(\mu) = - \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = -\mathbf{j}.$$

Hence we have

$$\iota(\rho) = P\rho P^{-1}, \quad P = \begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix} = \mathbf{i}.$$

In particular, we obtain the equation $\chi_\rho = \chi_{\iota(\rho)}$.

Next we show that an irreducible $\mathrm{SL}_2(\mathbb{C})$ -representation ρ satisfying $\chi_\rho = \chi_{\iota(\rho)}$ is metabelian. From the equations $\chi_{\iota(\rho)}(\mu) = -\chi_\rho(\mu)$ and $\chi_\rho = \chi_{\iota(\rho)}$, we have $\mathrm{tr} \rho(\mu) = 0$. Therefore by taking conjugation we can assume that $\rho(\mu)$ is given by the matrix $\mathbf{j}$.

We choose any two elements γ_1, γ_2 from $[\pi_1(\Sigma_K), \pi_1(\Sigma_K)]$ and set $\rho(\gamma_i)$ as

$$\rho(\gamma_i) = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix} \quad (i = 1, 2).$$

Since $\alpha(\gamma_i \mu) = t$, we have that $\chi_{\iota(\rho)}(\gamma_i \mu) = -\chi_\rho(\gamma_i \mu)$. From the assumption $\chi_\rho = \chi_{\iota(\rho)}$, we obtain $\mathrm{tr} \rho(\gamma_i \mu) = 0$. By Lemma 4.9, $b_i = c_i$ holds for $i = 1, 2$. Then the matrices $\rho(\gamma_1 \gamma_2)$ and $\rho(\gamma_1 \gamma_2)$ are given by

$$\begin{aligned} \rho(\gamma_1 \gamma_2) &= \begin{pmatrix} a_1 a_2 + b_1 b_2 & a_1 b_2 + b_1 d_2 \\ a_2 b_1 + b_2 d_1 & b_1 b_2 + d_1 d_2 \end{pmatrix}, \\ \rho(\gamma_2 \gamma_1) &= \begin{pmatrix} a_1 a_2 + b_1 b_2 & a_2 b_1 + b_2 d_1 \\ a_1 b_2 + b_1 d_2 & b_1 b_2 + d_1 d_2 \end{pmatrix}. \end{aligned}$$

Since $\alpha(\gamma_1\gamma_2\mu) = t$, we also have that $\text{tr } \rho(\gamma_1\gamma_2\mu) = 0$. By using Lemma 4.9 again, we have the equation:

$$a_1b_2 + b_1d_2 = a_2b_1 + b_2d_1.$$

Therefore $\rho(\gamma_1\gamma_2)$ is equal to $\rho(\gamma_2\gamma_1)$. The representation ρ is metabelian. $\square$ $\square$

By Proposition 4.4 and Proposition 4.8, we can conclude the following corollary.

Corollary 4.10.

$$X^{\text{irr}}(\Sigma_K)^{\widehat{\iota}} = \{\chi_\rho \mid \rho : \text{irreducible binary dihedral}\}.$$

The fixed point set $X(\Sigma_K)^{\widehat{\iota}}$ also could have characters of reducible representations. Actually, this fixed point set $X(\Sigma_K)^{\widehat{\iota}}$ contains only one character coming from reducible representations.

Before giving this proof, we define notations for the subsets in the character variety where the fixed point set $X(\Sigma_K)^{\widehat{\iota}}$ lives. Every character in such a subset send the meridian μ to the same value. We define such subsets as a level set of a function on the character variety.

Definition 4.11. For any given $\gamma \in \pi_1(\Sigma_K)$, we denote by I_γ a function on $X(\Sigma_K)$ defined by

$$\begin{aligned} I_\gamma : X(\Sigma_K) &\rightarrow \mathbb{C}, \\ \chi &\mapsto \chi(\gamma). \end{aligned}$$

This function is called the trace function of γ .

Definition 4.12. We set the subset $S_c(\Sigma_K)$ in $X(\Sigma_K)$ as the level set of I_μ at c , i.e.,

$$S_c(\Sigma_K) = I_\mu^{-1}(c).$$

This subset is called the slice at c .

We also set $R_c(\Sigma_K)$ as $R_c(\Sigma_K) := \{\rho \in R(\Sigma_K) \mid \text{tr } \rho(\mu) = c\}$.

Lemma 4.13. *The fixed point set $X(\Sigma_K)^{\widehat{\iota}}$ is contained in $S_0(\Sigma_K)$ and the restriction of $\widehat{\iota}$ gives an involution on $S_0(\Sigma_K)$.*

Proof. We have seen that every character χ in $X(\Sigma_K)^{\widehat{\iota}}$ satisfies that $\chi(\mu) = 0$ in the proof of Proposition 4.8. For any element χ in $S_0(\Sigma_K)$, we have

$$\widehat{\iota}(\chi)(\mu) = -\chi(\mu) = 0.$$

The image of $S_0(\Sigma_K)$ by $\widehat{\iota}$ is contained in $S_0(\Sigma_K)$ itself. Therefore the restriction of $\widehat{\iota}$ on $S_0(\Sigma_K)$ is an involution on $S_0(\Sigma_K)$. $\square$ $\square$

Remark 4.14. The involution $\widehat{\iota}$ (resp. ι) changes $S_c(\Sigma_K)$ (resp. $R_c(\Sigma_K)$) with $S_{-c}(\Sigma_K)$ (resp. $R_{-c}(\Sigma_K)$).

Every representation in $R_0(\Sigma_K)$ maps the meridian μ to matrices whose eigen values are $\pm\sqrt{-1}$. By Lemma 4.2 and $\Delta_K(-1) \neq 0$, we have no non-abelian reducible representation in $R_0(\Sigma_K)$.

Remark 4.15. The reducible part of $R_0(\Sigma_K)$ consists entirely of abelian representations.

Therefore, by Lemma 4.13, the set of reducible characters in $X(\Sigma_K)^{\widehat{\iota}}$ consists of only one character of the abelian representation $\varphi_{\sqrt{-1}}$. Summarizing, we have the following proposition.

Proposition 4.16. *The fixed point set $X(\Sigma_K)^\widehat{\iota}$ is expressed as*

$$X(\Sigma_K)^\widehat{\iota} = S_0(\Sigma_K)^\widehat{\iota} = \{\chi_\rho \mid \rho : \text{irreducible binary dihedral}\} \cup \{\chi_{\varphi_{\sqrt{-1}}}\}.$$

5. MAPS BETWEEN REPRESENTATION SPACES

In this section, we consider how to make an $\mathrm{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_n)$ from that of $\pi_1(\Sigma_K)$. To do this, we consider the maps between representation spaces defined by the pull-backs of the homomorphisms p_n , π and π_n between the fundamental groups. Then we have the following diagram among the $\mathrm{SL}_2(\mathbb{C})$ -representation spaces of $\pi_1(\Sigma_K)$, $\pi_1(C_n)$, $\pi_1(\Sigma)$ and $\pi_1(\Sigma_n)$:

$$\begin{array}{ccc} R(C_n) & \xleftarrow{\pi_n^*} & R(\Sigma_n) \\ p_n^* \uparrow & & \\ R(\Sigma_K) & \xleftarrow{\pi^*} & R(\Sigma). \end{array}$$

Note that π_n^* is injection since π_n is surjective. By Corollary 3.14, we can regard $\pi_n^*(R(\Sigma_n))$ as the set of representations in $R(C_n)$ such that every $\rho \in \pi_n^*(R(\Sigma_n))$ sends μ_n to the identity matrix. The intersection $p_n^*(R(\Sigma_K)) \cap \pi_n^*(R(\Sigma_n))$ in $R(C_n)$ gives a correspondence from $R(\Sigma_K)$ to $R(\Sigma_n)$. The intersection is given as follows.

Lemma 5.1.

$$(10) \quad p_n^*(R(\Sigma_K)) \cap \pi_n^*(R(\Sigma_n)) = \{p_n^*\rho \mid \rho \in R(\Sigma_K), \rho(\mu)^n = \mathbf{1}\}.$$

Proof. We choose an element ρ in $R(\Sigma_K)$ such that $p_n^*\rho$ is contained in $\mathrm{Im} \pi_n^*$. By Corollary 3.14 and $p(\mu_n) = \mu^n$, we have the following equivalent relations:

$$\begin{aligned} p_n^*\rho \in \mathrm{Im} \pi_n^* &\Leftrightarrow p_n^*\rho(\mu_n) = \mathbf{1} \\ &\Leftrightarrow \rho(\mu^n) = \mathbf{1}. \end{aligned}$$

□

□

Remark 5.2. The right hand side of Eq. (10) always has the following component

$$\{p_n^*\rho \mid \rho \in R(\Sigma_K), \rho(\mu) = \mathbf{1}\}.$$

By the isomorphism (6) and Corollary 3.14, we have the following identification:

$$\{\rho \in R(\Sigma_K) \mid \rho(\mu) = \mathbf{1}\} = R(\Sigma).$$

Through this identification, the restriction of the pull-back p_n^* coincides with the pull-back of the homomorphism $\pi_1(\Sigma_n) \rightarrow \pi_1(\Sigma)$.

From the intersection (10) in Lemma 5.1, we can make a correspondence from a subset in $R(\Sigma_K)$ into $R(\Sigma_n)$ as follows. For each $\gamma \in \pi_1(\Sigma_n)$, we let $\widehat{\gamma}$ be a lift to $\pi_1(C_n)$, i.e., $\pi_n(\widehat{\gamma}) = \gamma$. For a given $\rho \in R(\Sigma_K)$ satisfying $\rho(\mu)^n = \mathbf{1}$, an $\mathrm{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_n)$ is given by

$$(11) \quad \begin{aligned} &\pi_1(\Sigma_n) \rightarrow \mathrm{SL}_2(\mathbb{C}), \\ &\gamma \mapsto p_n^*\rho(\widehat{\gamma}). \end{aligned}$$

However if we consider the case $\Sigma = S^3$ and two-fold branched cover Σ_2 along a knot K , then the intersection $p_2^*(R(E_K)) \cap \pi_2^*(R(\Sigma_2))$ turns into

$$p_2^*R_2(\Sigma_K) \cup p_2^*R_{-2}(\Sigma_K) = \{p_2^*\rho \in R(C_2) \mid \rho : \pi_1(E_K) \rightarrow \{\pm \mathbf{1}\}\}.$$

This set has only two representations. Both of their restrictions to $\pi_1(\Sigma_2)$ are trivial. Hence we deal with only the slices $R_{\pm 2}(\Sigma_K)$ at ± 2 , which correspond to the second roots of unity, when we construct an $\mathrm{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_2)$ from that of $\pi_1(\Sigma_K)$.

In this section, we would like to find similar relationships between other slices and the character varieties of branched covers. Roughly speaking, our purpose in this section is to give a method to construct $\mathrm{SL}_2(\mathbb{C})$ -representations of $\pi_1(\Sigma_n)$ from not only slices $R_{2\cos(2k\pi/n)}(\Sigma_K)$ but also $R_{2\cos((2k-1)\pi/n)}(\Sigma_K)$. Hereafter we denote by $\xi(k/n)$ the number $2\cos(k\pi/n)$ simply. First we see that the pull-back of each element in the subset

$$(12) \quad \{\rho \in R(\Sigma_K) \mid \rho(\mu)^n = \pm 1\} \subset \bigcup_{k=1}^{n-1} R_{\xi(k/n)}(\Sigma_K) \cup \{\rho \in R(\Sigma_K) \mid \rho(\mu) = \pm 1\}$$

defines a $\mathrm{PSL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_n)$.

Lemma 5.3. *If ρ is an element in the subset (12), then the pull-back $p_n^*\rho$ defines $\mathrm{PSL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_n)$.*

Proof. We can regard $\pi_1(\Sigma_n)$ as the quotient $\pi_1(C_n)/|\langle \mu_n \rangle|$. It is enough to prove that $p_n^*\rho(\mu_n) = \pm 1$. This follows from direct calculations. $\square$ $\square$

Next we will lift these $\mathrm{PSL}_2(\mathbb{C})$ -representations of $\pi_1(\Sigma_n)$ to $\mathrm{SL}_2(\mathbb{C})$ by using a sign refinement in $R(C_n)$. As concerns the lifting problem on $\mathrm{PSL}_2(\mathbb{C})$ -representations to $\mathrm{SL}_2(\mathbb{C})$ -ones, several results have been known. For example, G. Burde [Bur90] has shown that binary dihedral representations form the branched point set of covering map from $\mathrm{SU}(2)$ -representations to $\mathrm{SO}(3)$ -representations for a two-bridge knot. Heusener and J. Porti [PH04] deal with the lifting problem for $\mathrm{PSL}_2(\mathbb{C})$ -representations of a finitely generated and presented group in context of the character variety.

Definition 5.4. Let $p_n^*\alpha : \pi_1(C_n) \rightarrow \langle t \rangle$ be the pull-back of α by p_n . Set $\zeta_m = \exp(2\pi\sqrt{-1}/m)$ and denote by $\alpha_{\zeta_{2n}}$ the composition of $p_n^*\alpha$ and an evaluation map $t = \zeta_{2n}$:

$$\begin{array}{ccc} \pi_1(C_n) & \xrightarrow{p_n^*\alpha} & \langle t^n \rangle \subset \langle t \rangle \\ & \searrow \alpha_{\zeta_{2n}} & \downarrow t=\zeta_{2n} \\ & & \{\pm 1\}. \end{array}$$

Then an involution ι_n on $R(C_n)$ is defined by

$$\begin{aligned} \iota_n : R(C_n) &\rightarrow R(C_n), \\ \rho &\mapsto \alpha_{\zeta_{2n}} \cdot \rho. \end{aligned}$$

This involution ι_n also induces an involution $\widehat{\iota}_n$ on the $\mathrm{SL}_2(\mathbb{C})$ -character variety $X(C_n)$ defined by $\widehat{\iota}_n(\chi_\rho) := \chi_{\iota_n(\rho)}$.

Now we construct a map Φ_n from the subset $\{\rho \in R(\Sigma_K) \mid \rho(\mu)^n = \pm 1\}$ into $R(\Sigma_n)$ as follows. Let ρ be an element of $R_{\xi(k/n)}(\Sigma_K)$ such that $\rho(\mu)^n = (-1)^k$. Then we define an $\mathrm{SL}_2(\mathbb{C})$ -representation $\Phi_n(\rho)$ of $\pi_1(C_n)$ by sending each element $\gamma \in \pi_1(C_n)$ to the following matrix:

$$\Phi_n(\rho)(\gamma) = \iota_n^k(p_n^*\rho)(\gamma) = (\alpha_{\zeta_{2n}}(\gamma))^k \cdot p_n^*\rho(\gamma),$$

where ι_n^k is k -times composition of ι_n .

Proposition 5.5. *For $\rho \in R_{\xi(k/n)}(\Sigma_K)$ satisfying $\rho(\mu)^n = (-\mathbf{1})^k$, we have the equality $\Phi_n(\rho)(\mu_n) = \mathbf{1}$. In particular, this homomorphism induces a homomorphism from $\pi_1(C_2)/|\langle \mu_n \rangle|$ into $\mathrm{SL}_2(\mathbb{C})$.*

Proof. This proposition follows from the following direct calculation. Let $\rho \in R_{\xi(k/n)}(\Sigma_K)$ satisfy $\rho(\mu)^n = (-\mathbf{1})^k$.

$$\begin{aligned}\Phi_n(\rho)(\mu_n) &= (\alpha_{\zeta_{2n}}(\mu_n))^k \cdot p_n^* \rho(\mu_n) \\ &= \zeta_{2n}^{kn} \cdot \rho(\mu)^n \\ &= (-1)^k (-\mathbf{1})^k \\ &= \mathbf{1}.\end{aligned}$$

By Corollary 3.14, this representation $\Phi_n(\rho)$ induces an $\mathrm{SL}_2(\mathbb{C})$ -representation of $\pi_1(C_n)/|\langle \mu_n \rangle| = \pi_1(\Sigma_n)$. $\square$ $\square$

The $\mathrm{SL}_2(\mathbb{C})$ -representation $\Phi_n(\rho)$ is a lift of the $\mathrm{PSL}_2(\mathbb{C})$ -representation $p_n^* \rho$ of $\pi_1(\Sigma_n)$:

$$\begin{array}{ccc} & & \mathrm{SL}_2(\mathbb{C}) \\ & \nearrow \Phi_n(\rho) & \downarrow \\ \pi_1(\Sigma_n) & \xrightarrow[p_n^* \rho]{} & \mathrm{PSL}_2(\mathbb{C}).\end{array}$$

Through the identification of $\pi_1(C_n)/|\langle \mu_n \rangle|$ with $\pi_1(\Sigma_n)$, we define an $\mathrm{SL}_2(\mathbb{C})$ -representation $\Phi_n(\rho)$ of $\pi_1(\Sigma_n)$. The domain $D(\Phi_n)$ of this map is given by

$$\begin{aligned}(13) \quad D(\Phi_n) &= \{\rho \in R(\Sigma_K) \mid \rho(\mu)^n = \pm \mathbf{1}\} \\ &= \bigcup_{k=1}^{n-1} \{\rho \in R_{\xi(k/n)}(\Sigma_K) \mid \rho(\mu)^n = \pm \mathbf{1}\} \cup \{\rho \in R(\Sigma_K) \mid \rho(\mu) = \pm \mathbf{1}\}.\end{aligned}$$

Remark 5.6. By Corollary 3.14, the set $\{\rho \in R(\Sigma_K) \mid \rho(\mu) = \mathbf{1}\}$ coincides with $\pi^*(R(\Sigma))$. Since the pull-back π^* is injective, we identify $\pi^*(R(\Sigma))$ with $R(\Sigma)$. The involution ι changes the two sets $\{\rho \in R(\Sigma_K) \mid \rho(\mu) = \pm \mathbf{1}\}$ each other. So we can rewrite the domain $D(\Phi_n)$ in (13) as

$$\bigcup_{k=1}^{n-1} \{\rho \in R_{\xi(k/n)}(\Sigma_K) \mid \rho(\mu)^n = \pm \mathbf{1}\} \cup R(\Sigma) \cup \iota(R(\Sigma)).$$

Summarizing, we obtain the following map between representations:

$$\Phi_n : D(\Phi_n) \rightarrow R(\Sigma_n),$$

where $\Phi_n(\rho)$ for a representation ρ in $R_{\xi(k/n)}(\Sigma_K)$ is given by $\Phi_n(\rho) = \iota_n^k(p_n^* \rho)$.

The correspondence Φ_n induces a map $\widehat{\Phi}_n$ on the subset $t(D(\Phi_n))$ in $X(\Sigma_K)$ defined by $\widehat{\Phi}_n(\chi_\rho) := \chi_{\Phi_n(\rho)}$. Then the domain $D(\widehat{\Phi}_n)$ of $\widehat{\Phi}_n$ is

$$(14) \quad D(\widehat{\Phi}_n) = \bigcup_{k=1}^{n-1} S_{\xi(k/n)}(\Sigma_K) \cup X(\Sigma) \cup \iota(X(\Sigma)) = \bigcup_{k=0}^n S_{\xi(k/n)}(\Sigma_K),$$

where we identify $\pi^*(X(\Sigma))$ with $X(\Sigma)$. Hence $\widehat{\Phi}_n$ is defined on $S_{\xi(k/n)}(\Sigma_K)$ as

$$(15) \quad \widehat{\Phi}_n : S_{\xi(k/n)}(\Sigma_K) \rightarrow X(\Sigma_n), \quad \chi_\rho \mapsto \chi_{\iota_n^k(p_n^* \rho)}.$$

By the definition of Φ_n , we can know what is the image of any abelian representation by Φ_n .

Lemma 5.7. *For every abelian representation ρ in $D(\Phi_n)$, the image $\Phi_n(\rho)$ is the trivial representation of $\pi_1(\Sigma_n)$.*

Proof. Since ρ is abelian, it sends all elements in the commutator subgroup of $\pi_1(\Sigma_K)$ to the identity matrix. For each generator $\tau_n^j(\tilde{x}_i)$ in the presentation of Lemma 3.15, the image by $\Phi(\rho)$ is expressed as $\Phi_n(\rho)(\tau_n^j(\tilde{x}_i)) = \rho(\mu^j x_i \mu^{-j})$. Since all x_i are contained in the commutator subgroup of $\pi_1(\Sigma_K)$, $\mu^j x_i \mu^{-j}$ is also an element in the commutator subgroup. Hence every $\mu^j x_i \mu^{-j}$ is sent to the identity matrix by ρ . Therefore $\Phi_n(\rho)$ is trivial. $\square$ $\square$

Remark 5.8. The involutions ι and $\hat{\iota}$ interchange slices as follows:

$$\iota(R_{\xi(k/n)}(\Sigma_K)) = R_{\xi(1-k/n)}(\Sigma_K), \quad \hat{\iota}(S_{\xi(k/n)}(\Sigma_K)) = S_{\xi(1-k/n)}(\Sigma_K).$$

We investigate the properties of the map Φ_n in the rest of this section. To consider the image of Φ_n , we introduce a notation for the action on $\pi_1(C_n)$ induced by the covering transformation.

First we define the following notion concerning τ_n to characterize the image of Φ_n .

Definition 5.9. An $\mathrm{SL}_2(\mathbb{C})$ -representation ρ of $\pi_1(C_n)$ is τ_n -equivariant if there exists an element C in $\mathrm{SL}_2(\mathbb{C})$ such that the following diagram is commutative:

$$(16) \quad \begin{array}{ccc} \pi_1(C_n) & \xrightarrow{\tau_n} & \pi_1(C_n) \\ \rho \downarrow & & \downarrow \rho \\ \mathrm{SL}_2(\mathbb{C}) & \xrightarrow{Ad_C} & \mathrm{SL}_2(\mathbb{C}), \end{array}$$

where $Ad_C(X) := CXC^{-1}$ for elements C and X in $\mathrm{SL}_2(\mathbb{C})$. We denote by $R^{\tau_n}(C_n)$ the set of τ_n -equivariant $\mathrm{SL}_2(\mathbb{C})$ -representations.

The τ_n -equivariance of ρ means that the pull-back of ρ by τ_n is expressed as the conjugation of an $\mathrm{SL}_2(\mathbb{C})$ -element.

Lemma 5.10. *If ρ is irreducible and τ_n -equivariant, then the matrix C is determined up to sign.*

Proof. It follows from Schur's lemma for ρ and $\tau_n^* \rho$. $\square$ $\square$

Similarly, we can also define τ_n -equivariant representations for $\pi_1(\Sigma_n, b_n)$. We denote by $R^{\tau_n}(\Sigma_n)$ the set of τ_n -equivariant representations of $\pi_1(\Sigma_n)$. We use the same symbol τ_n for the induced homomorphism on $\pi_1(\Sigma_n, b_n)$ by the covering transformation. The reason is due to the following Lemma.

Lemma 5.11. *The following diagram is commutative:*

$$(17) \quad \begin{array}{ccc} \pi_1(C_n) & \xrightarrow{\tau_n} & \pi_1(C_n) \\ \pi_n \downarrow & & \downarrow \pi_n \\ \pi_1(\Sigma_n) & \xrightarrow{\tau_n} & \pi_1(\Sigma_n). \end{array}$$

Proof. The fundamental group $\pi_1(\Sigma_n)$ is expressed as $\pi_1(C_n)/|\langle \mu_n \rangle|$. Since τ_n is automorphism and $\tau_n(\mu_n) = \mu_n$ in $\pi_1(C_n)$, together with the presentation of $|\langle \mu_n \rangle|$ as in Lemma 3.13, we have that $\tau_n(|\langle \mu_n \rangle|) = |\langle \mu_n \rangle|$. Hence τ_n induces an automorphism on the quotient group $\pi_1(C_n)/|\langle \mu_n \rangle|$. $\square$ $\square$

Moreover we can identify the set $\{\rho \in R^{\tau_n}(C_n) \mid \rho(\mu_n) = \mathbf{1}\}$ with $R^{\tau_n}(\Sigma_n)$ by the following Lemma 5.12.

Lemma 5.12. *The subset $\pi_n^*(R^{\tau_n}(\Sigma_n))$ coincides with $\{\rho \in R^{\tau_n}(C_n) \mid \rho(\mu_n) = \mathbf{1}\}$.*

Proof. This follows from the commutative diagram (17) in Lemma 5.11. $\square$ $\square$

The conjugation on representation spaces has the following property.

Lemma 5.13. *The conjugation for representations preserves τ_n -equivariance.*

Proof. Let ρ be a τ_n -equivariant representation of $\pi_1(C_n)$. There exists an element C in $\mathrm{SL}_2(\mathbb{C})$ such that the following diagram is commutative:

$$\begin{array}{ccc} \pi_1(C_n) & \xrightarrow{\tau_n} & \pi_1(C_n) \\ \rho \downarrow & & \downarrow \rho \\ \mathrm{SL}_2(\mathbb{C}) & \xrightarrow{Ad_C} & \mathrm{SL}_2(\mathbb{C}). \end{array}$$

Then for any P in $\mathrm{SL}_2(\mathbb{C})$, the representation $P\rho P^{-1}$ satisfies that

$$\begin{array}{ccc} \pi_1(C_n) & \xrightarrow{\tau_n} & \pi_1(C_n) \\ P\rho P^{-1} \downarrow & & \downarrow P\rho P^{-1} \\ \mathrm{SL}_2(\mathbb{C}) & \xrightarrow{Ad_{P\rho P^{-1}}} & \mathrm{SL}_2(\mathbb{C}). \end{array}$$

Hence $P\rho P^{-1}$ is τ_n -equivariant. $\square$ $\square$

In the following, we will prove that the image of Φ_n coincides with $R^{\tau_n}(\Sigma_n)$. We use the identification as in Lemma 5.12 and the following lemma.

$$\begin{aligned} R^{\tau_n}(\Sigma_n) &= \pi_n^*(R^{\tau_n}(\Sigma_n)) \\ &= \{\rho \in R^{\tau_n}(C_n) \mid \rho(\mu_n) = \mathbf{1}\}. \end{aligned}$$

Lemma 5.14. *Suppose that $\rho \in R^{\tau_n}(\Sigma_n)$ and C is a matrix as in the commutative diagram (16). The adjoint action Ad_{C^n} acts trivially on $\mathrm{Im} \rho$.*

Proof. The τ_n -equivariance of ρ gives us the equality that $\rho(\tau_n^i(\gamma)) = Ad_{c^i}(\rho(\gamma))$ for any γ in $\pi_1(C_n)$. For the case $i = n$, since $\tau_n^n(\gamma) = \mu_n \gamma \mu_n^{-1}$ and $\rho(\mu_n) = \mathbf{1}$, we have that $Ad_{C^n}(\rho(\gamma)) = \rho(\gamma)$ for any γ in $\pi_1(C_n)$, i.e., Ad_{C^n} acts trivially on $\mathrm{Im} \rho$. $\square$ $\square$

We next show the following property on p_n^* .

Lemma 5.15. *For every ρ in $R(\Sigma_K)$, the pull-back $p_n^*\rho$ is contained in $R^{\tau_n}(C_n)$.*

Proof. Let γ be an element in $\pi_1(C_n)$. By the following direct calculation, it follows that $p_n^*\rho$ is τ_n -equivariant:

$$\begin{aligned} p_n^*\rho(\tau_n(\gamma)) &= \rho(p_n(\tau_n(\gamma))) \\ &= \rho(\mu \cdot p_n(\gamma) \cdot \mu^{-1}) \\ &= \text{Ad}_{\rho(\mu)}(p_n^*\rho(\gamma)). \end{aligned}$$

□

□

Lemma 5.16. *We have the following inclusion*

$$\text{Im } \Phi_n \subset R^{\tau_n}(\Sigma_n),$$

where $R^{\tau_n}(\Sigma_n)$ is regarded as the subset in $R(C_n)$.

Proof. For any representation ρ in $D(\Phi_n) \cap R_{\xi(k/n)}(\Sigma_K)$ and each γ in $\pi_1(C_n)$, the matrix $\Phi_n(\rho)(\tau_n(\gamma))$ is given by

$$\begin{aligned} \Phi_n(\rho)(\tau_n(\gamma)) &= \alpha_{\zeta_{2n}}(\tau_n(\gamma))^k \cdot p_n^*\rho(\tau_n(\gamma)) \\ &= (\alpha_{\zeta_{2n}}(\gamma))^k \cdot \text{Ad}_{\rho(\mu)}(p_n^*\rho(\gamma)) \quad (\text{by Lemma 5.15}) \\ &= \text{Ad}_{\rho(\mu)}(\Phi_n(\rho)(\gamma)). \end{aligned}$$

Hence the $\text{SL}_2(\mathbb{C})$ -representation $\Phi_n(\rho)$ is τ_n -equivalent. □ □

In fact, the above inclusion gives an equality as sets.

Proposition 5.17. *The image of Φ_n coincides with the set of τ_n -equivariant $\text{SL}_2(\mathbb{C})$ -representations of $\pi_1(\Sigma_n)$, i.e.,*

$$\text{Im } \Phi_n = R^{\tau_n}(\Sigma_n).$$

Proof. By Lemma 5.16, it is enough to show the inclusion $\text{Im } \Phi_n \supset R^{\tau_n}(\Sigma_n)$. We regard $R^{\tau_n}(\Sigma_n)$ as a subset in $R^{\tau_n}(C_n)$. For each ρ in $R^{\tau_n}(\Sigma_n)$, we let C be a matrix as in the commutative diagram (16). We use a presentation of $\pi_1(C_n)$ induced from a Lin's presentation of $\pi_1(\Sigma_K)$ as in Lemma 3.10.

The relations of $\pi_1(C_n)$ give the equalities:

$$\rho(\mu_n \tilde{\alpha}_i^{(0)} \mu_n^{-1}) = \rho(\tilde{\beta}_i^{(n-1)}).$$

The left hand side turns into $\rho(\tilde{\alpha}_i^{(0)})$. The element $\tilde{\beta}_i^{(n-1)}$ can be written as $\tau_n^{n-1}(\tilde{\beta}_i^{(0)})$. Hence the right hand side $\rho(\tilde{\beta}_i^{(n-1)})$ turns into $\text{Ad}_{C^{n-1}}(\rho(\tilde{\beta}_i^{(0)}))$. By Lemma 5.14, Ad_{C^n} acts trivially on $\text{Im } \rho$. Hence we have the equalities

$$C\rho(\tilde{\alpha}_i^{(0)})C^{-1} = \rho(\tilde{\beta}_i^{(0)}).$$

To make a representation of $\pi_1(\Sigma_K)$, we set $\rho_0(\mu) = C$ and $\rho_0(x_i) = \rho(\tilde{x}_i)$. The relations $\rho_0(\mu\alpha_i\mu^{-1}) = C\rho(\tilde{\alpha}_i^{(0)})C^{-1}$ and $\rho_0(\beta_i) = \rho(\tilde{\beta}_i^{(0)})$ show that $\rho_0(\mu\alpha_i\mu^{-1}) = \rho_0(\beta_i)$ for all i . Hence ρ_0 is an $\text{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_K)$. We must show that ρ_0 is contained in $D(\Phi_n)$. By the following Lemma 5.18, there exists an integer k such that ρ_0 is contained in $R_{\xi(k/n)}(\Sigma_K)$. □ □

Lemma 5.18. *If ρ is an element in $R^{\tau_n}(\Sigma_n)$ and C is an $\text{SL}_2(\mathbb{C})$ -element as in the commutative diagram (16), then the matrix C^n is contained in $\{\pm \mathbf{1}\}$ or ρ is trivial.*

Proof. We regard ρ as an element in $R^{\tau_n}(C_n)$ satisfying $\rho(\mu_n) = \mathbf{1}$. By Lemma 5.14, the adjoint action Ad_{C^n} acts trivially on $\text{Im } \rho$, i.e., C^n commutes with all elements in $\text{Im } \rho \subset \text{SL}_2(\mathbb{C})$. If C and $\text{Im } \rho$ are not contained in the same maximal abelian subgroups in $\text{SL}_2(\mathbb{C})$, then C^n is contained in the intersection between the two different maximal subgroup (i.e., C^n is contained in $\{\pm \mathbf{1}\}$). In another case where C and $\text{Im } \rho$ are contained in the same maximal abelian subgroup, since in particular ρ is abelian, the representation ρ factors through the homology group $H_1(\Sigma_n; \mathbb{Z})$. By the presentation of $H_1(\Sigma_n; \mathbb{Z})$ as in Lemma 3.15, we have the following equality for the vectors of the generators in $H_1(\Sigma_n; \mathbb{Z})$.

$$\mathbf{x}_1 + \cdots + \mathbf{x}_{n-1} = \mathbf{0},$$

where

$$\mathbf{x}_i = {}^t([\tau_n^i(\tilde{x}_1)], \dots, [\tau_n^i(\tilde{x}_{2g})]).$$

This relation gives us the equality that

$$(18) \quad \rho(\tilde{x}_i) \rho(\tau_n(\tilde{x}_i)) \cdots \rho(\tau_n^{n-1}(\tilde{x}_i)) = \mathbf{1}.$$

Since ρ is abelian and τ_n -equivariant, we have that $\rho(\tau_n(\gamma)) = Ad_C(\rho(\gamma)) = \rho(\gamma)$ for any γ in $\pi_1(C_n)$. Together with Eq. (18), we can conclude that $\rho(\tilde{x}_i)^n = \mathbf{1}$. By taking conjugation, we can assume that for all $j = 1, \dots, n-1$,

$$\rho(\tau_n^j(\tilde{x}_i)) = \rho(\tilde{x}_i) = \begin{pmatrix} \zeta_n^{k_i} & 0 \\ 0 & \zeta_n^{-k_i} \end{pmatrix}.$$

On the other hand, from the relation $V\mathbf{x}_0 - U\mathbf{x}_{n-1} = \mathbf{0}$ in the presentation of $H_1(C_n; \mathbb{Z})$, we have the following equality:

$$(V - U) \begin{pmatrix} k_1 \\ \vdots \\ k_{2g} \end{pmatrix} = \mathbf{0} \pmod{\mathbb{Z}}.$$

By Remark 3.7, the determinant $\det(V - U)$ is $\Delta_K(1) = \pm 1$. So we obtain $\rho(\tilde{x}_i) = \mathbf{1}$. Since the homology classes of $\tau_n^j(\tilde{x}_i)$ form a basis of $H_1(\Sigma_n; \mathbb{Z})$, we conclude that the representation ρ is trivial. $\square$ $\square$

In the case of $\text{SU}(2)$ -representations, such a correspondence like Φ_n has been considered by Collin and Saveliev in [CS01]. They deal with irreducible $\text{SU}(2)$ -representations mainly. Here we focus on reducible representations of $\pi_1(\Sigma_n)$. Lin's presentations lead us to the property of Φ_n for reducible representations.

Now, Proposition 5.17 can be naturally extended to the set of characters as follows:

Proposition 5.19. *The image of $\widehat{\Phi}_n$ coincides with the fixed point set of the action τ_n^* on $X(\Sigma_n)$.*

Proof. The image $\text{Im } \widehat{\Phi}_n$ is given by $t(R^{\tau_n}(\Sigma_n))$. By definition, the subset $t(R^{\tau_n}(\Sigma_n))$ is contained in $X(\Sigma_n)^{\tau_n}$. We prove that every fixed point of τ_n^* in $X(\Sigma_n)$ is given by the character of a τ_n -equivariant representation.

First we consider the character $\chi_\rho \in X(\Sigma_n)^{\tau_n}$ associated to an irreducible representation ρ such that it is fixed by τ_n^* , i.e., $\tau_n^*(\chi_\rho) = \chi_{\tau_n^*\rho} = \chi_\rho$. Since ρ is irreducible, the representation ρ is conjugate to $\tau_n^*\rho$. This says that ρ is τ_n -equivariant.

Next we consider the case that $\chi_\rho \in X(\Sigma_n)^{\tau_n}$ is an reducible character such that $\tau_n^*(\chi_\rho) = \chi_\rho$. For every reducible representation there exists an abelian representation of $\pi_1(\Sigma_n)$ such that it has the same character as that of the reducible one and

the image is contained in the maximal abelian subgroup Hyp. So we can assume without loss of generality that $\text{Im } \rho$ is contained in Hyp. By using a presentation as in Lemma 3.15 and $\tau_n^*(\chi_\rho) = \chi_\rho$, we have the equality $\text{tr } \rho(\tau_n^j(\tilde{x}_i)) = \text{tr } \rho(\tilde{x}_i)$. Hence $\rho(\tau_n^j(\tilde{x}_i))$ is either $\rho(\tilde{x}_i)$ or $\rho(\tilde{x}_i)^{-1}$. Set $\rho(\tilde{x}_i)$ and $\rho(\tau_n^j(\tilde{x}_i))$ as

$$\rho(\tilde{x}_i) = \begin{pmatrix} r_i e^{\sqrt{-1}\theta_i} & 0 \\ 0 & r_i^{-1} e^{-\sqrt{-1}\theta_i} \end{pmatrix}, \quad \rho(\tau_n^j(\tilde{x}_i)) = \rho(\tilde{x}_i)^{\epsilon_j} \quad (\epsilon_j = \pm 1).$$

Since ρ factors through $H_1(\Sigma_n; \mathbb{Z})$, the relation $-U\mathbf{x}_j + V\mathbf{x}_{j+1}$ in $H_1(\Sigma_n; \mathbb{Z})$ gives us the following relation:

$$(19) \quad (-\epsilon_j U + \epsilon_{j+1} V) \begin{pmatrix} \log r_1 \\ \vdots \\ \log r_{2g} \end{pmatrix} = \mathbf{0}, \quad (-\epsilon_j U + \epsilon_{j+1} V) \begin{pmatrix} \theta_1 \\ \vdots \\ \theta_{2g} \end{pmatrix} \equiv \mathbf{0} \pmod{2\pi\mathbb{Z}}.$$

If there exists some j such that $\epsilon_{j+1} = \epsilon_j$, then it follows by the relation (19) and $\det(-U + V) = \Delta_K(1) = \pm 1$ that $\rho(\tilde{x}_i) = \mathbf{1}$ for all i , i.e., ρ is trivial. If the equality $\epsilon_{j+1} = -\epsilon_j$ holds for all j , then we have $\tau_n^* \rho = C \rho C^{-1}$ where $C = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Thus ρ is τ_n -equivariant in the both case. Therefore $\text{Im } \hat{\Phi}_n$ coincides with the fixed point set $X(\Sigma_n)^{\tau_n^*}$. $\square$ $\square$

We enclose this section with observation on the pre-images of reducible representation of $\pi_1(\Sigma_n)$ by Φ_n .

Proposition 5.20. *In the image of Φ_n , all reducible representations are abelian, i.e.,*

$$(20) \quad \text{Im } \Phi_n \cap R^{\text{red}}(\Sigma_n) = \text{Im } \Phi_n \cap R^{\text{ab}}(\Sigma_n).$$

Proof. We regard $R(\Sigma_n)$ as a subset in $R(C_n)$ by π_n^* . It is obvious that all abelian representations are contained in the set of reducible representations. We must show that the left hand side of Eq. (20) is contained in the right hand side.

By Proposition 5.17, the intersection of $\text{Im } \Phi_n$ with $R^{\text{red}}(\Sigma_n)$ is $R^{\tau_n}(\Sigma_n) \cap R^{\text{red}}(C_n)$ in $R(C_n)$. We suppose that ρ is an element in $R^{\text{red}}(C_n)$ such that $\rho(\mu_n) = \mathbf{1}$ and C is an $\text{SL}_2(\mathbb{C})$ -element satisfying $\tau_n^* \rho = \text{Ad}_C(\rho)$. We use the presentation of $\pi_1(C_n)$ as in Lemma 3.10. Since ρ is reducible, by taking conjugation, we can assume that

$$\rho(\tau_n^j(\tilde{x}_i)) = \begin{pmatrix} \alpha_{i,j} & \beta_{i,j} \\ 0 & \alpha_{i,j}^{-1} \end{pmatrix}$$

for all i and j . If $\alpha_{i,j}$ is ± 1 for all i and j , then $\text{Im } \rho$ is contained in the maximal abelian subgroup Para. Hence ρ is abelian. We consider the case that there exists a pair (i, j) such $\alpha_{i,j} \neq \pm 1$. If necessary, by taking conjugation of the following matrix

$$\begin{pmatrix} 1 & \beta_{i,j}/(\alpha_{i,j} - \alpha_{i,j}^{-1}) \\ 0 & 1 \end{pmatrix},$$

we can assume that $\rho(\tau_n^j(\tilde{x}_i))$ is a diagonal matrix. Since ρ is reducible, $\rho(\tau_n^{j+1}(\tilde{x}_i)) = C \rho(\tau_n^j(\tilde{x}_i)) C^{-1}$ is also an upper triangular matrix. Writing $C = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, we have that $cd = 0$. If c were zero, then we would have the following contradiction. When we set $\rho_0(\mu) = C$ and $\rho_0(x_i) = \rho(\tilde{x}_i)$, the correspondence ρ_0 defines an $\text{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_K)$. Moreover since the image of ρ_0 consists of upper

triangular matrices, the representation ρ_0 is reducible. Hence for every commutator γ in $\pi_1(\Sigma_K)$, the trace of $\rho_0(\gamma)$ is ± 2 . Applying this to the commutator x_i , we have that $\text{tr } \rho_0(x_i) = \alpha_{i,j} + \alpha_{i,j}^{-1} = \pm 2$. This is a contradiction to our assumption that $\alpha_{i,j} \neq \pm 1$. Therefore d is zero. Furthermore since $\rho(\tau_n^{j+h}(\tilde{x}_i)) = C^h \rho(\tau_n^j(\tilde{x}_i)) C^{-h}$ must be upper triangular, it follows that the matrix C turns into

$$\begin{pmatrix} 0 & b \\ b^{-1} & 0 \end{pmatrix}.$$

By reducibility of ρ , the matrix $\rho(\tau_n^k(\tilde{x}_l))$ is also sent to an upper triangular matrix by the conjugation of C . Therefore every $\beta_{k,l}$ must be zero. Then the image of ρ is contained in the maximal abelian subgroup Hyp, i.e., ρ is also abelian in this case. This completes the proof. $\square$ $\square$

Proposition 5.21. *The preimage $\Phi_n^{-1}(R^{\text{red}}(\Sigma_n))$ of reducible representations is expressed as*

$$\Phi_n^{-1}(R^{\text{red}}(\Sigma_n)) = R^{\text{meta}}(\Sigma_K) \cap D(\Phi_n).$$

Proof. By Proposition 5.20, it is enough to show that the preimage $\Phi_n^{-1}(R^{\text{ab}}(\Sigma_n))$ coincides with $R^{\text{meta}}(\Sigma_K) \cap D(\Phi_n)$. Let ρ be an element in $\Phi_n^{-1}(R^{\text{ab}}(\Sigma_n))$. By taking conjugation, we can assume that the image $\Phi_n(\rho)$ is contained in either the maximal abelian subgroups Hyp or Para. We use a Lin's presentation of $\pi_1(\Sigma_K)$ as in Lemma 3.3. In the case that $\text{Im } \Phi_n(\rho)$ is contained in Hyp, for all i the matrix $\rho(x_i)$ is expressed as

$$\rho(x_i) = \begin{pmatrix} \alpha_i & 0 \\ 0 & \alpha_i^{-1} \end{pmatrix}.$$

If all α_i are ± 1 , then ρ is abelian. We consider the case that there exist some α_i such that $\alpha_i \neq \pm 1$. Since $\Phi_n(\rho)(\tau_n(\tilde{x}_i)) = \pm \rho(\mu) \rho(x_i) \rho(\mu)^{-1}$ and this matrix is also upper triangular, the matrix $\rho(\mu)$ is expressed as either

$$\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 0 & b \\ -b^{-1} & 0 \end{pmatrix}.$$

In both cases, the representation ρ turns into metabelian.

In the case that $\text{Im } \Phi_n(\rho)$ is contained in Para, for all i , $\rho(x_i)$ is expressed as

$$\rho(x_i) = \pm \begin{pmatrix} 1 & \beta_i \\ 0 & 1 \end{pmatrix}.$$

If all β_i are zero, then ρ is abelian. We consider the case that there exists some non-zero β_i . Since $\Phi_n(\rho)(\tau_n(\tilde{x}_i)) = \pm \rho(\mu) \rho(x_i) \rho(\mu)$ is also a parabolic element, the matrix $\rho(\mu)$ must be upper triangular. Therefore ρ is reducible, in particular, it is metabelian.

On the other hand, by Remark 4.1 and the proof of Proposition 4.4, every metabelian representation is conjugate to either

$$\rho(x_i) = \begin{pmatrix} \alpha_i & 0 \\ 0 & \alpha_i^{-1} \end{pmatrix}, \quad \rho(\mu) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

or

$$\rho(x_i) = \begin{pmatrix} \pm 1 & \omega_i \\ 0 & \pm 1 \end{pmatrix}, \quad \rho(\mu) = \begin{pmatrix} \lambda & \beta \\ 0 & \lambda^{-1} \end{pmatrix}.$$

It follows from the definition of Φ_n that all metabelian representations in $D(\Phi_n)$ are sent to abelian ones of $\pi_1(\Sigma_n)$. $\square$ $\square$

6. ON THE GEOMETRY OF THE SLICE $S_0(\Sigma_K)$

In this section, we focus on the geometric/topological-like structure of the slice $S_0(E_K)$ for the exterior E_K of a knot K in S^3 . As described below, the slice $S_0(E_K)$ has a two-fold branched cover-like structure whose base space can be thought of as the $\mathrm{SL}_2(\mathbb{C})$ -character variety $X(\Sigma_2)$ and whose branched point set consists entirely of the characters of metabelian representations of the knot group. Note that the results in this section can be naturally extended to the case of an integral homology three-sphere with some modifications.

First we describe the representation spaces $R_0(E_K)$ and $R^{\tau_2}(\Sigma_2)$ in detail.

Lemma 6.1. *If $\rho \in R_0(E_K)$, then $\rho(\mu)^2 = -1$. Therefore the domain of Φ_2 is given by the union $R_0(E_K) \cup R(S^3) \cup \iota(R(S^3)) = R_0(E_K) \cup \{\text{trivial rep.}\} \cup \{\iota\}$. Moreover the image of $\widehat{\Phi}_2$ coincides with that of the restriction on $R_0(E_K)$.*

Proof. This follows from the Cayley–Hamilton identity and Remark 5.6. The equality $\mathrm{Im} \widehat{\Phi}_2 = \mathrm{Im} \widehat{\Phi}_2|_{R_0(E_K)}$ can be deduced from the fact that $R(S^3)$ and $\iota(R(S^3))$ consists of only the trivial representation and ι , respectively. It follows from Lemma 5.7 that they are sent to the trivial representation of $\pi_1(\Sigma_2)$ by Φ_2 and all abelian representations in $R_0(E_K)$ are also sent to the trivial one of $\pi_1(\Sigma_2)$. This shows that $\mathrm{Im} \widehat{\Phi}_2 = \mathrm{Im} \widehat{\Phi}_2|_{R_0(E_K)}$. □ □

By Lemma 6.1, in the case of knots in S^3 , the domain $D(\Phi_2)$ is reduced to the slice $R_0(E_K)$. This makes the properties on Φ_2 and $R_0(E_K)$ much clear. For example, we obtain the next proposition by Proposition 5.17, Lemma 5.7, Propositions 5.20 and 5.21.

Proposition 6.2. *The map Φ_2 from $R_0(E_K)$ onto $R^{\tau_2}(\Sigma_2)$ makes the following correspondence. For an element ρ in $R_0(E_K)$,*

- (1) *If ρ is abelian, then $\Phi_2(\rho)$ is the trivial representation of $\pi_1(\Sigma_2)$;*
- (2) *If ρ is irreducible and metabelian, then $\Phi_2(\rho)$ is a non-trivial abelian representation in $R^{\tau_2}(\Sigma_2)$; and*
- (3) *If ρ is non-metabelian, in particular irreducible, then $\Phi_2(\rho)$ is an irreducible representation in $R^{\tau_2}(\Sigma_2)$.*

Recall that the map Φ_2 induces the map $\widehat{\Phi}_2 : S_0(E_K) \rightarrow X(\Sigma_2)$ defined by $\widehat{\Phi}_2(\chi_\rho) := \chi_{\Phi_2(\rho)}$ (see also (15)). This gives us the following theorem.

Theorem 6.3. *The image of $S_0(E_K)$ by $\widehat{\Phi}_2$ coincides with the subset $X(\Sigma_2)^{\tau_2^*} = \{\chi_\rho \in X(\Sigma_2) \mid \rho : \tau_2\text{-equivariant}\}$. The map $\widehat{\Phi}_2 : S_0(E_K) \rightarrow X(\Sigma_2)^{\tau_2^*}$ is 1 : 1-correspondence on the fixed point set $S_0(E_K)^{\widehat{\iota}}$ and 2 : 1-correspondence on $S_0(E_K) \setminus S_0(E_K)^{\widehat{\iota}}$. Moreover two points in the preimage of a point in $X(\Sigma_2)^{\tau_2^*}$ are changed each other by the involution $\widehat{\iota}$ on $S_0(E_K)$.*

This theorem gives the two-fold branched cover-like structure of the slice $S_0(K)$, which is one of the main results in this article. We would like to show Theorem 6.3 in the rest of this section.

Lemma 6.4. *If ρ is an element in $D(\Phi_2)$, then the map Φ_2 sends both ρ and $\iota(\rho)$ to the same $\mathrm{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_2)$, i.e., $\Phi_2(\rho) = \Phi_2(\iota(\rho))$. Moreover, this equality gives $\widehat{\Phi}_2(\chi_\rho) = \widehat{\Phi}_2(\chi_{\iota(\rho)})$.*

Proof. The map Φ_2 is constructed by $\alpha_{\sqrt{-1}}$ and the pull-back p_2^* . For each element ρ in $R_0(E_K)$, the $\mathrm{SL}_2(\mathbb{C})$ -representation $\alpha_{\sqrt{-1}} \cdot p^*(\iota(\rho))$ is expressed as follows:

$$\begin{aligned}\alpha_{\sqrt{-1}} \cdot p_2^*(\iota(\rho)) &= \alpha_{\sqrt{-1}} \cdot p_2^*(\alpha_{-1} \cdot \rho) \\ &= \alpha_{-\sqrt{-1}} \cdot p_2^* \rho \\ &= \alpha_{\sqrt{-1}} \cdot p_2^* \rho.\end{aligned}$$

Hence $\Phi_2(\iota(\rho))$ coincides with $\Phi_2(\rho)$. This equality also holds for the induced map $\widehat{\Phi}_2$. $\square$ $\square$

By Proposition 4.8, the fixed point set $S_0(E_K)^\wedge$ in $S_0(E_K)$ consists entirely of the characters of metabelian representations, which are in $R_0(E_K)$.

Proposition 6.5. *The restriction of $\widehat{\Phi}_2$ on $S_0(E_K)^\wedge$ gives a bijection between $S_0(E_K)^\wedge$ and all characters of abelian representations of $\pi_1(\Sigma_2)$.*

Proof. By Proposition 4.8 and Lemma 4.13, we focus on the characters of metabelian representations. By Propositions 5.20 and 5.21, the map $\widehat{\Phi}_2$ sends the characters of metabelian representations onto the characters of abelian ones of $\pi_1(\Sigma_2)$. By Propositions 4.4 and 4.16, the set of characters of metabelian representations consists of $((|\Delta_K(-1)| - 1)/2 + 1)$ characters, where $\Delta_K(t)$ is the Alexander polynomial of K . On the other hand, we see that the set of characters of abelian representations of $\pi_1(\Sigma_2)$ also consists of $((|\Delta_K(-1)| - 1)/2 + 1)$ characters as follows.

Every abelian representation factors through the abelianization $H_1(\Sigma_2; \mathbb{Z})$. Since the order of $H_1(\Sigma_2; \mathbb{Z})$ is finite (an odd integer $|\Delta_K(-1)|$), $H_1(\Sigma_2; \mathbb{Z})$ is decomposed into the direct sum of some finite cyclic groups. The character of an abelian representation is determined by the traces for generators of these cyclic groups. By conjugation, the images of generators of cyclic groups are given by the diagonal matrices whose diagonal components are roots of unity (because the order of each cyclic group is finite). Combining these facts, we can show that the number of characters of abelian representations is given by $(|\Delta_K(-1)| - 1)/2 + 1$. Therefore the restriction of $\widehat{\Phi}_2$ on $S_0(E_K)^\wedge$ gives a one-to-one correspondence. $\square$ $\square$

Proposition 6.6. *The restriction of $\widehat{\Phi}_2$ on $S_0(E_K) \setminus S_0(E_K)^\wedge$ gives a 2 : 1-correspondence.*

Proof. It is sufficient to show that if ρ and ρ' are two non-metabelian representations of $\pi_1(E_K)$ satisfying $\widehat{\Phi}_2(\chi_\rho) = \widehat{\Phi}_2(\chi_{\rho'})$, then $\chi_{\rho'}$ coincides with either χ_ρ or $\bar{\iota}(\chi_\rho)$. By Proposition 6.2, the representations $\Phi_2(\rho)$ and $\Phi_2(\rho')$ are irreducible. Since $\Phi_2(\rho)$ and $\Phi_2(\rho')$ have the same characters. By [CS83, Proposition 1.5.2], these two representations are conjugate, i.e., there exists an $\mathrm{SL}_2(\mathbb{C})$ -element P such that

$$(21) \quad \Phi_2(\rho') = P\Phi_2(\rho)P^{-1}.$$

We use the presentations of $\pi_1(E_K)$ and $\pi_1(\Sigma_2)$ as in Lemma 3.15. By the construction of Φ_2 , the matrix $\Phi_2(\rho)(\tilde{x}_i)$ is expressed as

$$(22) \quad \Phi_2(\rho)(\tilde{x}_i) = \rho(x_i),$$

Combining Eq. (22) with Eq. (21), we have the equality

$$(23) \quad \rho'(x_i) = P\rho(x_i)P^{-1}.$$

Similarly, the matrix $\Phi_2(\rho)(\tau_2(\tilde{x}_i))$ is expressed as

$$(24) \quad \Phi_2(\rho)(\tau_2(\tilde{x}_i)) = \rho(\mu)\rho(x_i)\rho(\mu)^{-1} = \mathrm{Ad}_{\rho(\mu)}(\Phi_2(\rho)(\tilde{x}_i)).$$

Combining Eq. (24) with Eq. (21), we have the equality

$$\Phi_2(\rho')(\tau_2(\tilde{x}_i)) = Ad_{P\rho(\mu)P^{-1}}(\Phi_2(\rho')(\tilde{x}_i)).$$

The matrix $\Phi_2(\rho)(\tau_2(\tilde{x}_i))$ is also expressed as $Ad_{\rho'(\mu)}(\Phi_2(\rho')(\tilde{x}_i))$. By Lemma 5.10, we obtain the equality

$$(25) \quad \rho'(\mu) = \pm P\rho(\mu)P^{-1}.$$

From the relations that $\iota(\rho)(x_i) = \rho(x_i)$ and $\iota(\rho)(\mu) = -\rho(\mu)$ and Eqs. (23) and (25), we can conclude that ρ' is conjugate to either ρ or $\iota(\rho)$ by P . $\square$ $\square$

Proof of Theorem 6.3. Combining Lemma 6.4, Proposition 6.5 and Proposition 6.6, we obtain Theorem 6.3. $\square$ $\square$

Now, we observe that the set of τ_2 -equivariant representations $R^{\tau_2}(\Sigma_2)$ contains the set of all abelian representations $R^{\text{ab}}(\Sigma_2)$.

Lemma 6.7. *The set $R^{\tau_2}(\Sigma_2)$ contains $R^{\text{ab}}(\Sigma_2)$.*

Proof. Let ρ be an abelian $\text{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma_2)$. Since this representation factors through $H_1(\Sigma_2; \mathbb{Z})$ which is a finite abelian group with the order $|\Delta_K(-1)|$. By taking conjugation, the image $\text{Im } \rho$ is contained in the maximal abelian subgroup Hyp . So we can set $\rho(\tilde{x}_i)$ as

$$\rho(\tilde{x}_i) = \begin{pmatrix} \alpha_i & 0 \\ 0 & \alpha_i^{-1} \end{pmatrix}.$$

By Lemma 3.15, we have $[\tilde{x}_i] + [\tau_2(\tilde{x}_i)] = \mathbf{0}$ in $H_1(\Sigma_2; \mathbb{Z})$ for all i . Hence $\rho(\tau_2(\tilde{x}_i))$ is given by $\rho(\tilde{x}_i)^{-1}$.

If we set $C = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, then the diagram (16) becomes commutative. Therefore ρ is τ_2 -equivariant. $\square$ $\square$

Remark 6.8. Lemma 6.7 does not hold for three-fold branched covers of S^3 . As shown below, for the trefoil knot in S^3 there exists an abelian representation of $\pi_1(\Sigma_3)$ which is not τ_3 -equivariant.

Proposition 6.9. *Let K be the left hand trefoil knot in S^3 . The set $R^{\text{ab}}(\Sigma_3)$ is not contained in $R^{\tau_3}(\Sigma_3)$.*

To show Proposition 6.9, we consider the diagram of the left hand trefoil and the Heegaard splitting associated to the Seifert surface depicted in Figure 4. We fix a spine as in Figure 4. The Seifert matrix is given by the following matrix

$$Q = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}.$$

We decompose S^3 into two handle bodies as in Figure 5. Each attaching circle of two-handles intersects with the boundaries of meridian circles of one-handles at only one point in this Heegaard splitting. The Morse complex $C_*^{\text{Morse}}(S^3; \mathbb{Z})$ is obtained by the Heegaard splitting. The boundary operator $\partial_2 : C_2^{\text{Morse}}(S^3; \mathbb{Z}) \rightarrow C_1^{\text{Morse}}(S^3; \mathbb{Z})$ is expressed as the identity matrix. The relations in Lin's presentation are given by

$$\alpha_1 = x_1 x_2^{-1}, \quad \alpha_2 = x_2, \quad \beta_1 = x_1, \quad \beta_2 = x_1^{-1} x_2.$$

Hence, by Lemma 3.15, a presentation of $\pi_1(\Sigma_3)$ is given by

$$\langle \tilde{x}_1, \tilde{x}_2, \tau_3(\tilde{x}_1), \tau_3(\tilde{x}_2), \tau_3^2(\tilde{x}_1), \tau_3^2(\tilde{x}_2), |\tilde{\alpha}_1^{(k)} = \tilde{\beta}_1^{(k-1)}, \tilde{\alpha}_2^{(k)} = \tilde{\beta}_2^{(k-1)} (k \bmod 3) \rangle.$$

This presentation is reduced as

$$\langle \tilde{x}_1, \tilde{x}_2 | \tilde{x}_1 \tilde{x}_2^{-1} = \tilde{x}_2 \tilde{x}_1, \tilde{x}_2 \tilde{x}_1 = \tilde{x}_1^{-1} \tilde{x}_2 \rangle.$$

About another method to obtain this presentation, for example see [Rol90, Chapter 10].




FIGURE 4. The Seifert surface of the trefoil knot




FIGURE 5. Heegaard splitting of S^3 associated to S

Proof of Proposition 6.9. By Lemma 3.15, the homology group $H_1(\Sigma_3; \mathbb{Z})$ is presented as

$$\mathbb{Z}^6 \xrightarrow{A} \mathbb{Z}^6 \rightarrow H_1(\Sigma_3; \mathbb{Z}),$$

where

$$A = \begin{pmatrix} {}^tQ & -Q & 0 \\ 0 & {}^tQ & -Q \\ -Q & 0 & {}^tQ \end{pmatrix}.$$

If we set

$$\begin{cases} \rho(\tilde{x}_1) = \mathbf{1}, & \rho(\tau_3(\tilde{x}_1)) = -\mathbf{1}, & \rho(\tau_3^2(\tilde{x}_1)) = -\mathbf{1}, \\ \rho(\tilde{x}_2) = -\mathbf{1}, & \rho(\tau_3(\tilde{x}_2)) = \mathbf{1}, & \rho(\tau_3^2(\tilde{x}_2)) = -\mathbf{1}, \end{cases}$$

then this correspondence defines an abelian representation of $\pi_1(\Sigma_3)$. Since the trace of $\rho(\tilde{x}_1)$ is not equal to that of $\rho(\tau_3(\tilde{x}_1))$, there does not exist any $\mathrm{SL}_2(\mathbb{C})$ -element such that the diagram (16) commutes. $\square$ $\square$

7. APPLICATIONS

In the previous section, we have investigated the correspondence $\widehat{\Phi}_2$ between $S_0(E_K)$ and $X(\Sigma_2)^{\tau_2^*}$. In particular, Lemma 6.7 shows that $R^{\text{ab}}(\Sigma_2) \subset R^{\tau_2}(\Sigma_2)$. In this section, we look into surjectivity of $\widehat{\Phi}_2 : S_0(E_K) \rightarrow X(\Sigma_2)$ for two-bridge knots and pretzel knots of type (p, q, r) . Namely, we show that $R^{\tau_2}(\Sigma_2)$ contains all $\text{SL}_2(\mathbb{C})$ -representations $R(\Sigma_2)$.

As regards two-bridge knots, it is well-known that the two-fold branched cover Σ_2 along a two-bridge knot is a three-dimensional lens space. Since the fundamental group of a lens space is cyclic, the $\text{SL}_2(\mathbb{C})$ -representation space for a lens space consists entirely of abelian representations. By Theorem 6.3 and Proposition 6.2, the slice $S_0(E_K)$ consists of characters of metabelian representations of $\pi_1(E_K)$. It is known that the order of $H_1(\Sigma_2; \mathbb{Z})$ is given by $|\Delta_K(-1)|$. Thus we have proved the following Lemma, which follows originally from the proof of Theorem 1.3 in [Nag07].

Lemma 7.1. *For a two-bridge knot K , the slice $S_0(E_K)$ coincides with the fixed point set $S_0(E_K)^{\widehat{\tau}}$ and it can be identified with the set of $((|\Delta_K(-1)| - 1)/2 + 1)$ conjugacy classes of the $\text{SL}_2(\mathbb{C})$ -metabelian representations of $\pi_1(E_K)$.*

Another way of stating Lemma 7.1 is to say that if Σ_2 is a lens space, then the slice $S_0(E_K)$ coincides with the fixed point set $S_0(E_K)^{\widehat{\tau}}$. This means that the difference $S_0(E_K) \setminus S_0(E_K)^{\widehat{\tau}}$ is an obstruction for Σ_2 to be a lens space.

Theorem 7.2. *If $S_0(E_K) \setminus S_0(E_K)^{\widehat{\tau}} \neq \emptyset$, then Σ_2 is not a lens space.*

Remark 7.3. The similar statement also holds for a knot in an integral homology three-sphere.

Here we give explicit representatives in conjugacy classes of metabelian representations by using R. Riley's construction and the correspondence Φ_2 . We will show the following theorem.

Theorem 7.4. *Let K be a two-bridge knot and p the determinant of K , i.e., $p = |\Delta_K(-1)|$. We assume that a presentation of $\pi_1(E_K)$ is given by $\langle x, y \mid wx = yw \rangle$ where w is a word in x and y . Then the slice $S_0(E_K)$ is identified with the following set of conjugacy classes of $\text{SL}_2(\mathbb{C})$ -representations:*

$$\{[\rho_k] \mid k = 1, \dots, (p-1)/2\} \cup \{[\varphi_{\sqrt{-1}}]\},$$

where the representation ρ_K is given by

$$x \mapsto \begin{pmatrix} \sqrt{-1} & -\sqrt{-1} \\ 0 & -\sqrt{-1} \end{pmatrix}, \quad y \mapsto \begin{pmatrix} \sqrt{-1} & 0 \\ -\sqrt{-1}(e^{k\pi\sqrt{-1}/p} - e^{-k\pi\sqrt{-1}/p})^2 & -\sqrt{-1} \end{pmatrix}$$

and $\varphi_{\sqrt{-1}}$ is the abelian representation such that

$$\varphi_{\sqrt{-1}}(\mu) = \begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix}.$$

Riley had shown a construction of non-abelian $\text{SL}_2(\mathbb{C})$ -representations of two-bridge knot groups in [Ril84] by using a polynomial equation $\phi_K(t, u) = 0$. His construction gives every representative in each conjugacy class of a non-abelian representation. Note that the irreducible metabelian representations are given by using roots of $\phi_K(-1, u) = 0$. We also obtain the next statement about $\phi_K(-1, u)$.

Theorem 7.5. *We keep notations in Theorem 7.4. The polynomial $\phi_K(-1, u)$ has distinct $(p-1)/2$ roots $\{(e^{k\pi\sqrt{-1}/p} - e^{-k\pi\sqrt{-1}/p})^2 \mid k = 1, \dots, (p-1)/2\}$. Namely, $\phi_K(-1, u)$ is expressed as*

$$\phi_K(-1, u) = (-1)^{(p-1)/2} \prod_{k=1}^{(p-1)/2} \left\{ u - (e^{k\pi\sqrt{-1}/p} - e^{-k\pi\sqrt{-1}/p})^2 \right\}.$$

Note that a generator of $\pi_1(\Sigma_2)$ in S^3 is illustrated as in Figure 6 (for details about the generator, see [BZ03, Chapter 12]).




FIGURE 6. A generator γ of $\pi_1(\Sigma_2)$

Proof of Theorem 7.4. Let K be a two-bridge knot. We choose the following presentation of $\pi_1(E_K)$ obtained by reducing a Wirtinger presentation as

$$\langle x, y \mid wx = yw \rangle,$$

where the generators x and y are given in Figure 6 and w is a word in x and y . We focus on the non-abelian part in $S_0(E_K)$. By Theorem 1 in [Ril84], every conjugacy class of non-abelian $\mathrm{SL}_2(\mathbb{C})$ -representations has representatives given by the following representation ρ :

$$\rho(x) = \begin{pmatrix} \sqrt{t} & 1/\sqrt{t} \\ 0 & 1/\sqrt{t} \end{pmatrix}, \quad \rho(y) = \begin{pmatrix} \sqrt{t} & 0 \\ -u\sqrt{t} & 1/\sqrt{t} \end{pmatrix},$$

where t and u satisfy the equation that $\phi_K(t, u) = 0$. We now consider the case that $t = -1$ to describe elements in $S_0(E_K)$. A generator γ of $\pi_1(\Sigma_2)$ is given by Figure 6. If we regard γ as in E_K , then γ is expressed as $\gamma = xy^{-1}$. The image of γ by the abelianization homomorphism α is 1. Then $\Phi_2(\rho)(\gamma)$ is expressed as follows:

$$\begin{aligned} \Phi_2(\rho)(\gamma) &= \hat{\alpha}_{\sqrt{-1}}(\gamma) \cdot \rho(p(\gamma)) \\ &= \rho(xy^{-1}) \\ &= \begin{pmatrix} u+1 & 1 \\ u & 1 \end{pmatrix}. \end{aligned}$$

By Theorem 6.3, The representation $\Phi_2(\rho)$ is abelian. Every abelian representation of $\pi_1(\Sigma_2)$ is determined by the eigen values for the generator γ . Since $\pi_1(\Sigma_2)$ is the

cyclic group with order p . Therefore the matrix $\Phi_2(\rho)(\gamma)$ is conjugate to

$$\begin{pmatrix} e^{2\pi\sqrt{-1}/p} & 0 \\ 0 & e^{-2\pi\sqrt{-1}/p} \end{pmatrix}.$$

Comparing the traces of $\Phi_2(\rho)(\gamma)$ and the above diagonal matrix, we have that

$$u = (e^{\pi\sqrt{-1}/p} - e^{-\pi\sqrt{-1}/p})^2.$$

Since Φ_2 gives a one-to-one correspondence on characters of metabelian representations, we can conclude the statement. $\square$ $\square$

Proof of Theorem 7.5. If we fix a choice of square root of -1 , the roots of $\phi_K(-1, u)$ correspond to the representatives of non-abelian part in $S_0(E_K)$ by one-to-one. From [Ril84], the highest degree of u in $\phi_K(t, u)$ is given by the determinant of K , i.e., $p = |\Delta_K(-1)|$. Moreover it follows from the proof of Proposition 3 in [Ril84] that the coefficient of leading term of $\phi_K(t, u)$ is equal to $(-1)^{(p-1)/2}$. By the degree of u and Theorem 7.4, all roots of $\phi_K(-1, u)$ are given precisely by $(p-1)/2$ distinct real numbers:

$$\{(e^{k\pi\sqrt{-1}/p} - e^{-k\pi\sqrt{-1}/p})^2 \mid k = 1, \dots, (p-1)/2\}.$$

Therefore we obtained the equation:

$$\phi_K(-1, u) = (-1)^{(p-1)/2} \prod_{k=1}^{(p-1)/2} \left\{ u - (e^{k\pi\sqrt{-1}/p} - e^{-k\pi\sqrt{-1}/p})^2 \right\}.$$

$\square$

$\square$

Example. Let K be the trefoil knot. Then $\phi_K(t, u)$ is given by $u - t - t^{-1} + 1$. The root of $\phi_K(-1, u)$ is $u = -3$. The corresponding representation ρ is expressed as

$$\rho(x) = \begin{pmatrix} \sqrt{-1} & -\sqrt{-1} \\ 0 & -\sqrt{-1} \end{pmatrix}, \quad \rho(y) = \begin{pmatrix} \sqrt{-1} & 0 \\ 3\sqrt{-1} & -\sqrt{-1} \end{pmatrix}.$$

Then the matrix $\Phi_2(\rho)(xy^{-1})$ is given by

$$\begin{pmatrix} -2 & 1 \\ -3 & 1 \end{pmatrix}.$$

In fact, we have that $\Phi_2(\rho)(xy^{-1})^3 = \mathbf{1}$ by direct calculations. Since $|\Delta_K(-1)| = 3$, the formula in Theorem 7.5 turns into

$$u - (e^{\pi\sqrt{-1}/3} - e^{-\pi\sqrt{-1}/3})^2 = u + 3.$$

This polynomial coincides with $\phi_K(-1, u)$.

We remark that the map Φ_2 is not injective in general. For example, for any pretzel knot Φ_2 is not injective while it is surjective.

Proposition 7.6. *If K is a pretzel knot of type (p, q, r) , then the map $\hat{\Phi}_2$ from $S_0(E_K)$ to $X(\Sigma_2)$ is surjective.*

The key to Proposition 7.6 is that the two-fold branched cover Σ_2 along a pretzel knot of type (p, q, r) is the Brieskorn manifold of type (p, q, r) . The fundamental group of this Brieskorn manifold has a presentation with four generators s_1, s_2, s_3 and h (a central element) as that of a Seifert manifold (for more details, see [BZ03, Chapter 12] and [Sav02]). In fact, it is more convenient to work with another set of generators $t_1 = s_1, t_2 = s_1 s_2$ and h when we consider the induced action on

the fundamental group by the covering transformation. We use this idea to prove Proposition 7.6.

Proof. The character variety $X(\Sigma_2)$ is expressed as the union $X^{\text{ab}}(\Sigma_2) \cup X^{\text{irr}}(\Sigma_2)$. We will show that each part is contained in $\text{Im } \widehat{\Phi}_2$. By Proposition 6.5, $X^{\text{ab}}(\Sigma_2)$ is contained in $\text{Im } \widehat{\Phi}_2$. Hence it suffices to show that $X^{\text{irr}}(\Sigma_2)$ is contained in $\text{Im } \widehat{\Phi}_2$. Since $\text{Im } \widehat{\Phi}$ is $X(\Sigma_2)^{\tau_2^*}$, we check the inclusion $X^{\text{irr}}(\Sigma_2) \subset X(\Sigma_2)^{\tau_2^*}$. For an irreducible character χ_ρ , we have the following equivalence relations:

$$\begin{aligned} \tau_2^*(\chi_\rho) = \chi_\rho &\Leftrightarrow \chi_{\tau_2^*\rho} = \chi_\rho \\ &\Leftrightarrow \rho \stackrel{\text{conj}}{\sim} \tau_2^*\rho \\ &\Leftrightarrow \rho \in R^{\tau_2}(\Sigma_2). \end{aligned}$$

So, to complete the proof, it is enough to show that every irreducible representation is τ_2 -equivariant. The fundamental group $\pi_1(\Sigma_2)$ has the following presentation:

$$\langle s_1, s_2, s_3, h \mid s_1^p h = 1, s_2^q h = 1, s_3^r h = 1, [s_i, h] = 1 \ (1 \leq i \leq 3), s_1 s_2 s_3 = 1 \rangle.$$

For the generators in this presentations, the action of τ_2 is expressed as follows:

$$\tau_2 : h \mapsto h, s_1 \mapsto s_1^{-1}, s_2 \mapsto s_1 s_2^{-1} s_1^{-1}, s_3 \mapsto s_1 s_2 s_3^{-1} s_2^{-1} s_1^{-1}.$$

We set t_1 and t_2 as $t_1 = s_1$ and $t_2 = s_1 s_2$. Then three elements h, t_1 and t_2 generate the group $\pi_1(\Sigma_2)$. For the new generators, the action of τ_2 is expressed as

$$\tau_2 : h \mapsto h, t_1 \mapsto t_1^{-1}, t_2 \mapsto t_2^{-1}.$$

Since ρ is irreducible, the image $\text{Im } \rho$ is a non-abelian subgroup in $\text{SL}_2(\mathbb{C})$. Hence the central element $\rho(h)$ in $\text{Im } \rho$ is $\pm \mathbf{1}$. By the relation $t_1^p h$, the matrix $\rho(t_1)$ has a finite order, in particular it is hyperbolic. By taking conjugation, we can suppose that the irreducible representation ρ is given by

$$\rho(h) = \pm \mathbf{1}, \quad \rho(t_1) = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}, \quad \rho(t_2) = \begin{pmatrix} s & t \\ u & v \end{pmatrix},$$

where $ut \neq 0$. We set a complex number γ as $\gamma^2 = \sqrt{-u/t}$. Moreover, taking a conjugate by the matrix

$$\begin{pmatrix} \gamma & 0 \\ 0 & \gamma^{-1} \end{pmatrix} \quad (\gamma^2 = \sqrt{-u/t}),$$

we can assume that $\rho(t_2)$ is given by

$$\begin{pmatrix} s & t \\ -t & v \end{pmatrix}$$

at the beginning.

For a unit quaternion $\mathbf{k} = \begin{pmatrix} 0 & \sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix}$, the action of conjugation of $\mathbf{k}$ is expressed as

$$\mathbf{k} \rho(h) \mathbf{k}^{-1} = \rho(h), \quad \mathbf{k} \rho(t_1) \mathbf{k}^{-1} = \rho(t_1)^{-1}, \quad \mathbf{k} \rho(t_2) \mathbf{k}^{-1} = \rho(t_2)^{-1}.$$

Therefore it follows that $\tau_2^* \rho$ coincides with $\mathbf{k} \rho \mathbf{k}^{-1}$, i.e., ρ is τ_2 -equivariant.

Together with $X^{\text{ab}}(\Sigma_2) \subset X(\Sigma_2)^{\tau_2^*}$, we have that $X(\Sigma_2) \subset X(\Sigma)^{\tau_2} = \text{Im } \widehat{\Phi}_2$, namely $\widehat{\Phi}_2$ is surjective. □

ACKNOWLEDGMENTS

The authors would like to thank Professor Xiao-Song Lin for giving substantial comments which drew their attention to the notion of the metabelian representations when the first author stayed at the University of California, Riverside in 2006. The first author had been partially supported by JSPS Research Fellowships for Young Scientists and has been partially supported by the Grant-in-Aid for Young Scientists (Start-up). The second author would like to thank Professor Toshitake Kohno for his advice and support. The second author had been partially supported by the 21st century COE program at Graduate School of Mathematical Sciences, University of Tokyo.

REFERENCES

- [AM90] S. Akbulut and J. D. McCarthy. *Casson's invariant for oriented homology 3-spheres: an exposition*. Mathematical Notes **36**. Princeton University Press, 1990.
- [BC06] H. Boden and C. Curtis. The $SL_2(\mathbb{C})$ casson invariant for seifert fibered homology spheres and surgeries on twist knots. *J. Knot Theory Ramifications*, **15**:813–837, 2006.
- [Bur67] G. Burde. Darstellungen von knotengruppen. *Math. Ann.*, **173**:24–33, 1967.
- [Bur90] G. Burde. $SU(2)$ -representation spaces for two-bridge knot groups. *Math. Ann.*, **288**:103–119, 1990.
- [BZ03] G. Burde and H. Zieschang. *Knots*. Walter de Gruyter & Co., Berlin, 2003.
- [CS83] M. Culler and P. Shalen. Varieties of group representations and splittings of 3-manifolds. *Ann. of Math.*, **117**:109–146, 1983.
- [CS01] O. Collin and N. Saveliev. Equivariant Casson invariants via gauge theory. *J. Reine Angew. Math.*, **541**:143–169, 2001.
- [Cur01] C. Curtis. An intersection theory count of the $SL_2(\mathbb{C})$ -representations of the fundamental group of a 3-manifold. *Topology*, **40**:773–787, 2001.
- [dR67] G. de Rham. Introduction aux polynômes d'un nœud. *Enseignement Math.*(2), **13**:187–194, 1967.
- [Her97] C. Herald. Flat connections, the Alexander invariant, and Casson's invariant. *Comm. Anal. Geom.*, **5**:93–120, 1997.
- [HK98] M. Heusener and J. Kroll. Deforming abelian $SU(2)$ -representations of knot groups. *Comment. Math. Helv.*, **73**:480–498, 1998.
- [HPS01] M. Heusener, J. Porti, and E. Suárez. Deformations of reducible representations of 3-manifold groups into $SL_2(\mathbb{C})$. *J. Reine Angew. Math.*, **530**:191–227, 2001.
- [Lic97] W. B. R. Lickorish. *An introduction to knot theory*. Graduate Texts in Mathematics **175**. Springer-Verlag, New York, 1997.
- [Lin92] X.-S. Lin. A knot invariant via representation spaces. *J. Differential Geom.*, **35**:337–357, 1992.
- [Lin01] X.-S. Lin. Representations of knot groups and twisted Alexander polynomials. *Acta Math. Sin. (Engl. Ser.)*, **17**:361–380, 2001.
- [Nag07] F. Nagasato. Finiteness of a section of the $SL(2, \mathbb{C})$ -character variety of knot groups. *to appear in Kobe Journal of Mathematics*, **24**, 2007.
- [OS04] P. Ozsváth and Z. Szabó. Holomorphic disks and topological invariants for closed three-manifolds. *Ann. of Math. (2)*, **159**:1027–1158, 2004.
- [PH04] J. Porti and M. Heusener. The variety of characters in $PSL(2, \mathbb{C})$. *Bol. Soc. Mat. Mex.*, **10**:221–237, 2004.
- [Ril84] R. Riley. Nonabelian representations of 2-bridge knot groups. *Quart. J. Math. Oxford Ser. (2)*, **35**:191–208, 1984.
- [Rol90] D. Rolfsen. *Knots and links*. Publish or Perish Inc., 1990.
- [Sav99] N. Saveliev. *Lectures on the topology of 3-manifolds*. de Gruyter Textbook. Walter de Gruyter & Co., Berlin, 1999.
- [Sav02] N. Saveliev. Representation spaces of seifert fibered homology spheres. *Topology and its Applications*, **126**:49–61, 2002.

DEPARTMENT OF MATHEMATICS, MEIJO UNIVERSITY, SHIOGAMAGUCHI TEMPAKU, NAGOYA 468-8502, JAPAN

E-mail address: `fukky@ccmfs.meijo-u.ac.jp`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1 KOMABA MEGURO, TOKYO 153-8914, JAPAN

E-mail address: `shouji@ms.u-tokyo.ac.jp`